



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

April 29, 2016

Joseph L. Powell
Projects Manager
Valent U.S.A. Corporation
1600 Rivera Avenue, Suite 200
Walnut Creek, CA 94596-8025

Subject: PRIA Label Amendment –Adding new minor uses for Onion, bulb subgroup 03-07A, Vegetable, fruiting group 08-10, Fruit, pome group 11-10, Fruit, stone group 12-12, Berry, low growing, subgroup 13-07G, except cranberry, Rapeseed subgroup 20A, except flax, Sunflower subgroup 20B, Cotton Seed subgroup 20C, and Stevia; Adding supplemental label for new uses
Product Name: V-10137 1 EC Herbicide
EPA Registration Number: 59639-132
Application Date: 10/10/2014
Decision Numbers: 497969 & 497968

Dear Mr. Powell:

The application referred to above, submitted under the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable under FIFRA sec 3 (c)(5). You must submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

This supplemental labeling contains some new and/or revised uses and/or directions which may be additional to the uses and/or directions found on the label on or attached to the container, but this supplemental labeling does not by itself constitute the complete set of use directions. The complete set of use directions is set forth on the container label as combined with this supplemental labeling.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40


CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Lisa Pahel by phone at (703) 347-0459, or via email at pahel.lisa@epa.gov.



Rachel C. Holloman, Chief
Fungicide and Herbicide Branch,
Registration Division, (7505P)
Office of Pesticide Programs

Enclosure



GROUP 1 HERBICIDE

[Bracketed text is optional]

V-10137 1 EC HERBICIDE

Active Ingredient	By Wt
*Clethodim.....	12.6%
Other Ingredients.....	87.4%
Total	100.0%

*(E)-2-[1-[[[(3-chloro-2-propenyl)oxy]imino]propyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one

Contains Petroleum Distillates

V-10137 1 EC Herbicide is an emulsifiable concentrate containing 0.97 lb clethodim active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE NEXT PAGE FOR ADDITIONAL PRECAUTIONARY STATEMENTS

**NET CONTENTS 2.5 Gallon
15 Gallon
120 Gallon**

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Harmful if swallowed.

FIRST AID	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Immediately call a poison control center or doctor.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give any liquid to the person.• Do not give anything to an unconscious person.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-892-0099 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Ingestion of this product or subsequent vomiting can result in aspiration of light hydrocarbon liquid, which can cause pneumonitis. If ingested, probable mucosal damage may contraindicate the use of gastric lavage.	

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves such as barrier laminate or viton \geq 14 mils, shoes plus socks and protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove clothing/PPE immediately after handling this product. Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS:

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwater or rinsate.

The use of this product may pose a hazard to the federally designated endangered species of Solano Grass and Wild Rice. Use of this product is prohibited in the following areas where the species are known to exist:

Solano Grass: Solano County, California: the vernal lakes area bounded by the Union Pacific Railroad and Hastings Road to the north, Highway 113 to the east, Highway 12 to the south and Travis Air Force Base to the west.

Wild Rice: Hays County, Texas.

PHYSICAL OR CHEMICAL HAZARDS:

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL AND PAMPHLET. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls over short-sleeved shirt and short pants, chemical-resistant gloves such as barrier laminate or viton \geq 14 mils and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep all unprotected persons out of operating areas, or vicinity where there may be drift.
Do not enter treated areas without protective clothing until sprays have dried.

**DISCLAIMER, RISKS OF USING THIS PRODUCT,
LIMITED WARRANTY
AND LIMITATION OF LIABILITY**

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE FULLEST EXTENT ALLOWED BY LAW AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label **and subject to the Risks of Using This Product as described above. To the extent consistent with applicable law AND AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED.** No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

To the fullest extent allowed by law Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. TO THE FULLEST EXTENT ALLOWED BY LAW THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

To the extent consistent with applicable law allowing such requirements Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is latter, so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law if Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing **Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability**, which may not be modified by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, to the extent allowed by applicable law. **Always read and follow the restrictions and limitations for all products whether used alone or in a tank mix. The most restrictive labeling of any product used applies in tank mixtures, including all crop rotational and other crop restrictions.**

TABLE OF CONTENTS

CROP INFORMATION	
Restrictions	
Precautions	
Control Symptoms	
APPLICATION INFORMATION	
Timing of Applications.....	
Ground Application	
Air Application.....	
Spot Treatment.....	
Chemigation – Garlic and Onion (Dry Bulb and Green) Sprinkler Irrigation Application.....	
Restrictions.....	
Use Precautions.....	
SPRAY DRIFT MANAGEMENT	
Restrictions.....	
RESISTANCE MANAGEMENT	

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-10137 1 EC HERBICIDE.....

Use Directions for Annual Grasses (All Crops).....	
Use Directions for Annual & Perennial Grass Control in Established Alfalfa and Mint with V-10137 1 EC Herbicide.....	
Directions For Reduced Rate Use To Control Small Annual Grasses Canola [including Rapeseed Subgroup 20A Except Flax Seed, Mustard Seed and Sesame Seed; Flax; Legume Vegetables (Dry and Succulent); Mustard Seed; Soybean; Sugar Beet and Sunflower Table	
Use Directions for Perennial Grasses (All Crops) Table.....	
Use Directions for Annual Bluegrass Control with V-10137 1 EC Herbicide All Crops	
Use Directions for Use in Roundup Ready® Field Corn (Burndown)	

TANK MIX - LABEL INFORMATION.....

Tank Mix Application of V-10137 1 EC Herbicide and Broadleaf Herbicides for Control of Grasses and Broadleaf Weeds	
Mixing Instructions.....	
Information on Antagonism	

Alfalfa

Table 2. V-10137 1 EC Herbicide Tank Mixes with Broadleaf Herbicides for Alfalfa.....	
---	--

Use Directions for Bushberry, Caneberry, Pome Fruit and Stone Fruit.....

Canola [Including Rapeseed Subgroup 20A, Except Flax Seed, Mustard Seed and Sesame Seed]	
---	--

Table 3. Reduced Rate V-10137 1 EC Herbicide Tank Mixes with Broadleaf Herbicides	
---	--

Cotton

Table 4. V-10137 1 EC Herbicide Tank Mixed with Cobra and MSMA applied Post Directed to Cotton.....	
Table 5. V-10137 1 EC Herbicide Tank Mixed with Glyphosate to Control Emerged Grasses in Cotton as a Broadcast Application	

Dry and Succulent Shelled Beans

Table 6. V-10137 1 EC Herbicide Tank Mixes with Broadleaf Herbicides for Dry and Succulent Shelled Beans	
---	--

Flax

Table 7. Reduced Rate V-10137 1 EC Herbicide Tank Mixes with Broadleaf Herbicides for Flax.....	
---	--

Peanut

Table 8. V-10137 1 EC Herbicide Tank Mixes with Broadleaf Herbicides For Peanut.....	
Use Directions for Grass Suppression for Harvest Efficiency in Peanut with V-10137 1 EC Herbicide.....	

Soybean

Table 9. V-10137 1 EC Herbicide Tank Mix with Broadleaf Herbicides for the Control of Volunteer Corn (Including Roundup Ready) in Soybean.....	
Table 10. V-10137 1 EC Herbicide Tank Mixes with Broadleaf Herbicides for Soybean.....	

Sugar Beet

Table 11. V-10137 1 EC Herbicide Tank Mixed with Broadleaf Sugar Beet Herbicides.....	
Table 12. V-10137 1 EC Herbicide Plus Betanex or Betamix Tank Mix for Three Sequential Applications for Annual Grass Control (Micro-Rate Application)	
Directions for Use for Micro-Rate Applications to Sugar Beet.....	
Table 13. Tank Mix Application of V-10137 1 EC Herbicide and Fungicides for Control of Grass Weeds and Diseases in Sugar Beet.....	
Table 14. Tank Mix Application of V-10137 1 EC Herbicide and Insecticides for Control of Grass Weeds and Insects in Alfalfa, Cotton, Mint, Peanut (including perennial), Soybean and Sunflower	

Fallow Land

Directions for Use	
Precautions.....	
Restrictions.....	
Table 15. V-10137 1 EC Herbicide in Tank Mixes to Control Annual and Perennial Grasses in Fallow Land.....	
Use Directions for Grass Suppression in Non-Crop Areas with V-10137 1 EC Herbicide	
Table 16. V-10137 1 EC Herbicide for the Control and/or Suppression of Tall Fescue in Native Prairie Warm-Season Grass Restoration Projects	
Special Application Instructions/Precautions	
Table 17. V-10137 1 EC Herbicide for the Suppression of Tall Fescue Seed-Heads in Non-Producing Agricultural Areas	
Special Application Instructions/Precautions	

Directions For Use in Ornamentals

Ornamental Trees	
Ground Covers.....	
Garden Flowers and Plants.....	
Shrubs.....	
Use Directions for Annual Grasses in Ornamentals.....	
Use Directions for Annual Bluegrass Control with V-10137 1 EC Herbicide in Ornamentals.....	
Use Directions for Perennial Grasses in Ornamentals.....	
Non-Bearing Food Crops	
Conifer Trees	
Non-Crop or Non-Planted Areas	
Storage and Disposal.....	

CROP INFORMATION

FOR USE ON: Alfalfa, Artichoke (Globe), Asparagus, Bean (dry) and Pea (shelled)[Dry shelled Bean and Pea (except Soybean Subgroup 6C)]¹, Bean and Pea (succulent shelled) [Succulent shelled Pea and Bean Subgroup 6B]², Berry Low Growing (except Cranberry and Strawberry [Subgroup 13-07G]³, Broccoli, Bushberry⁴, Cabbage, Canola*⁵, Caneberry [Subgroup 13-07A]⁶, Carrot, Cauliflower (and other Head and Stem Brassica Vegetables) [Subgroup 5A]⁷, Celery, Clover (grown in Idaho, Oregon and Washington only), Conifers, Cotton (including cotton grown for seed), Cranberry, Cucumber, Eggplant (and other Fruiting Vegetables [Crop Group 8-10])⁸, Fallow Land (and other non-producing agricultural areas), Field Corn⁹, Flax*, Garden Beet, Garlic, Herbs [Subgroup 19A]¹⁰, Hops, Horseradish (and other Root Vegetables [Subgroup 1A])¹¹, Legume Vegetables (edible podded) [Subgroup 6A]¹², Lettuce, Head and Leaf (and other Leafy Greens [Subgroup 4A])¹³, Melons (including Cantaloupe and Watermelon)¹⁴, Mint, Mustard Greens (and other Leafy Brassica Greens [Subgroup 5B])¹⁵ Mustard Seed*, Non-Bearing Food Crops, Non-Crop or Non-Planted Areas, Onions (dry bulb) [Subgroup 3-07A]¹⁶, Onion (green), Ornamentals, Peanut (including perennial), Peppers (bell and non-bell), Pome Fruit [Crop Group 11-10]¹⁷, Potato, Radish, Rhubarb (and other Leaf Petioles) [Subgroup 4B]¹⁸, Safflower, Sesame, Squash (including Pumpkin)¹⁴, Soybean, Stevia (dried leaves), Stone Fruit [Crop Group 12-12]¹⁹, Strawberry, Sugar Beet, Sunflower [Subgroup 20B]²⁰, Sweet Potato, Turnip Greens, Tomato and Yam (and other Tuberous and Corm Vegetables) [Subgroup 1D]²¹

*Not for use in California

- 1 Other Bean (dry) and Pea (shelled) [Dry Shelled Bean and Pea except Soybean Subgroup 6C] crops approved for use with V-10137 1 EC Herbicide include: Bean (*Lupinus* spp.), grain lupin, sweet lupin, white lupin and white sweet lupin; Bean (*Phaseolus* spp.), field, kidney, lima (dry), navy, pinto and tepary; Bean (*Vigna* spp.), adzuki bean, black-eyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, broad (dry), chickpea (garbanzo), guar, lablab bean and lentil; Pea (*Pisum* spp.), field and pigeon.
- 2 Other Bean & Pea (succulent shelled) [Subgroup 6B] crops approve for use with V-10137 1 EC Herbicide include: Bean (*Phaseolus* spp.), broad bean (succulent), lima bean (green); Bean (*Vigna* spp.), black-eyed pea, cowpea, Southern pea; Pea (*Pisum* spp.), English pea, garden pea, green pea and pigeon pea.
- 3 Other Berry Low Growing(except Cranberry and Strawberry) [Subgroup 13-07G] crops approved for use with V-10137 1 EC Herbicide include: Bearberry, Bilberry, Blueberry (lowbush), Cloudberry, Lingonberry, Muntries and Partridgeberry.
- 4 Other Bushberry crops approved for use with V-10137 1 EC Herbicide include: aronia berry, blueberry (highbush), black currant, buffalo currant, Chilean guava, cranberry (highbush), elderberry, European barberry, gooseberry, honeysuckle (edible), huckleberry (edible), jostaberry, juneberry (saskatoon berry), native currant, salal, sea buckthorn, red currant; cultivars, varieties and/or hybrids of these.
- 5 Other canola [Including Rapeseed Subgroup 20A (except flax seed, mustard seed and sesame seed)] crops approved for use with V-10137 1 EC Herbicide include: borage, crambe, cuphea, echium, gold of pleasure (camelina),hare's ear mustard, lesquerella, lunaria, meadowfoam, milkweed, oil radish, poppy seed, rapeseed (canola), sweet rocket
- 6 Other Caneberry [Subgroup 13-07A] crops approved for use with V-10137 1 EC Herbicide include: Blackberry, Loganberry, Raspberry (black and red), Wild Raspberry, cultivars, varieties and/or hybrids of these.
- 7 Other head and stem brassica vegetables [Subgroup 5A] crops approved for use with V-10137 1 EC Herbicide include: Chinese broccoli, Brussels sprouts, Chinese (napa) cabbage, Chinese mustard, cavalo broccolo and kohlrabi.

- 8 Other Fruiting Vegetables (except tomato) [Crop Group 8-10] crops approved for use with V-10137 1 EC Herbicide include: African eggplant; bush tomato; bell pepper; cocona; currant tomato; eggplant; garden huckleberry; goji berry; groundcherry; martynia; naranjilla; okra; pea eggplant; pepino, nonbell pepper; roselle; scarlet eggplant; sunberry; tomatillo; tree tomato; cultivars, varieties, and/or hybrids of these.
- 9 For burndown of existing stand of Roundup Ready field corn or volunteer Roundup Ready field corn prior to replanting field corn.
- 10 Other Herb [Subgroup 19A] crops approved for use with V-10137 1 EC Herbicide include: angelica, balm, basil, borage, burnet, camomile, catnip, chervil (dried), chive, Chinese chive, clary, coriander (leaf), costmary, culantro (leaf), curry (leaf), dill (dillweed), horehound, hyssop, lavender, lovage (leaf), marigold, marjoram (*Origanum* spp.), nasturtium, parsley (dried), pennyroyal, rosemary, rue, sage and savory, summer and winter.
- 11 Other root vegetables [Subgroup 1A] crops approved for use with V-10137 1 EC Herbicide include: burdock, edible; celeriac; chervil, turnip-rooted; chicory; ginseng; parsley, turnip-rooted; parsnip; radish, oriental; rutabaga; salsify; salsify, black; salsify, Spanish; skirret and turnip.
- 12 Other Edible Podded Legume Vegetable [Subgroup 6A] crops approved for use with V-10137 1 EC Herbicide include: Bean (*Phaseolus* spp.), runner, snap and wax; Bean (*Vigna* spp.), asparagus, Chinese longbean, moth, yardlong, jackbean; Pea (*Pisum* spp.), dwarf, edible-pod, snow, sugar snap, pigeon and sword bean.
- 13 Other Leafy Greens [Subgroup 4A] crops approved for use with V-10137 1 EC Herbicide include: amaranth (Chinese spinach, leafy amaranth and tampala), arugula (rocket), chervil, chrysanthemum (edible-leaved and garland), corn salad, cress (garden, yellow rock and winter), dandelion, dock (sorrel), endive (escarole), lettuce (head and leaf), orach, parsley, purslane (garden and winter), radicchio (red chicory), spinach, spinach (New Zealand and Vine (Indian and Malabar)).
- 14 Other cucurbit vegetables [Crop Group 9] crops approved for use with V-10137 1 EC Herbicide include: Chayote (fruit), Chinese Wax Gourd, Citron Melon, Edible Gourd, Gherkin and Muskmelons (all) including Honeydew Melon.
- 15 Other leafy brassica greens [Subgroup 5B] crops approved for use with V-10137 1 EC Herbicide include: broccoli raab, Chinese (bok choy) cabbage, collards, kale, mizuna, mustard greens, mustard spinach, rape greens.
- 16 Other Onion, bulb [Subgroup 3-07A] crops approved for use with V-10137 1 EC Herbicide include: Daylily, bulb; fritillaria, bulb; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; lily, bulb; onion, bulb; onion, Chinese, bulb; onion, pearl; onion, potato, bulb; shallot, bulb; cultivars, varieties, and/or hybrids of these.
- 17 Other pome fruit [Crop Group 11-10] crops approved for use with V-10137 1 EC Herbicide include: apple, azarole, crabapple, loquat, mayhaw, medlar, pear, Asian pear, quince, Chinese quince, Japanese quince and tejocote.
- 18 Other leaf petiole crops approved for use with V-10137 1 EC Herbicide include: cardoon, celtuce, Chinese celery, Florence fennel, and Swiss chard.
- 19 Other stone fruit [Crop Group 12-12] crops approved for use with V-10137 1 EC Herbicide include: apricot; , Japanese apricot, capulin, black cherry, nanking cherry, sweet cherry, tart cherry, Chinese jujube, nectarine; peach; plum; American plum, beach plum, Canada plum, cherry plum, chickasaw plum, damson plum, Japanese plum, Klamath plum, prune plum, plumcot, sloe
- 20 Other sunflower [Subgroup 20B] crops approved for use with V-10137 1 EC Herbicide include: calendula, castor oil plant, Chinese tallowtree, euphorbia, evening primrose, jojoba, Niger seed, rose hip, stokes aster, tallowwood, tea oil plant, vernoia
- 21 Other tuberous and corm vegetables [Subgroup 1D] crops approved for use with V-10137 1 EC Herbicide include: arracacha, arrowroot, Chinese artichoke, Jerusalem artichoke, edible canna, bitter and sweet cassava, chayote (root), chufa, dasheen (taro), ginger, leren, tanier, turmeric; yam, bean and yam, true.

V-10137 1 EC Herbicide is a selective postemergence herbicide for control of annual and perennial grasses. V-10137 1 EC Herbicide does not control sedges or broadleaf weeds.

RESTRICTIONS

- Application on Long Island, New York, is restricted to no more than 32 fl oz of V-10137 1 EC Herbicide (0.25 lb ai) per acre per year.
- Do not apply if rain is expected within 1 hour of application, as control may be unsatisfactory.
- Do not plant rotational crops until 30 days after application of V-10137 1 EC Herbicide unless crop is listed on V-10137 1 EC Herbicide label.
- Do not apply a postemergence broadleaf herbicide within one day following application of V-10137 1 EC Herbicide or reduced grass control may result.
- Do not apply under conditions of stress. Applying V-10137 1 EC Herbicide under conditions that do not promote active grass growth will reduce herbicide effectiveness. These conditions include drought, excessive water, extremes in temperature, low humidity and grasses either partially controlled or stunted from prior pesticide applications. Grasses under these kinds of stressful conditions will not absorb and translocate V-10137 1 EC Herbicide effectively, and will be less susceptible to herbicide activity.

PRECAUTIONS

- Grass crops such as corn, rice, small grains, sorghum or turf, etc. are highly sensitive to *V-10137 1 EC* Herbicide.
- *V-10137 1 EC* Herbicide is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided.
- Optimal perennial grass control can be obtained if rhizomes or stolons are cut up by preplant tillage practices (discing, plowing, etc.) to stimulate maximum emergence of grass shoots. Cultural practices, such as continuous no-tillage in which the perennial grass rhizomes or stolons are not cut up, results in a very staggered, non-uniform weed emergence. Due to this non-uniform weed emergence, no fewer than two *V-10137 1 EC* Herbicide applications per season per year are recommended at the appropriate weed-growth stage rate under continuous no-till conditions.
- While all the vegetable crops on this label have been tested and are tolerant to *V-10137 1 EC* Herbicide, not all specialty varieties of these crops have been tested. It is advised that, before applying *V-10137 1 EC* Herbicide to specialty varieties of vegetable crops on this label, crop tolerance be investigated first using a small section of the field. It is possible that injury symptoms can occur. Symptoms may appear as leaf speckling or stunting.
- Always read and follow the restrictions and limitations for all products whether used alone or in a tank mix. The most restrictive labeling of any product used applies in tank mixtures, including all crop rotational and other crop restrictions.
- Tank mixes of *V-10137 1 EC* Herbicide and broadleaf herbicides may result in reduced grass control. If grass regrowth occurs, an additional application of *V-10137 1 EC* Herbicide may be necessary.

Control Symptoms

Treated grass weeds show a reduction in vigor and growth. Early chlorosis/necrosis of younger plant tissue is followed by a progressive collapse of the remaining foliage. Symptoms will generally be observed in 7 to 14 days after application, depending on grass species treated and environmental conditions.

APPLICATION INFORMATION

Timing of Applications

Apply *V-10137 1 EC* Herbicide postemergence to actively growing grasses according to prescribed rates in the tables. Applications made to grass plants stressed by insufficient moisture, hot or cold temperatures, or to grass plants exceeding recommended growth stages may result in unsatisfactory control. **Restriction:** Do not apply under these conditions.

In arid regions where irrigation is used to supplement limited rainfall, *V-10137 1 EC* Herbicide should be applied, as soon as possible, after an irrigation (within 7 days). In arid regions, a second application of *V-10137 1 EC* Herbicide will generally provide more effective control of perennial grass weeds than a single application. Make second application to actively growing grass 2 to 3 weeks after emergence of new growth.

Cultivation of treated grasses 7 days prior to or within 7 days after application of *V-10137 1 EC* Herbicide may reduce weed control.

Ground Application

Use of sufficient spray volumes and pressure is essential to ensure complete coverage. Use a minimum of 5 gals and a maximum of 40 gals of spray solution per acre. Under the following conditions a minimum of 10 gals per acre is required: ultra narrow row cotton, narrow row soybeans, broadleaf herbicide tank mixes, perennial grasses, volunteer corn, drought or stress conditions, heavy grass pressure or when grasses are at or near maximum height. Failure to use a minimum of 10 gals per acre under these conditions can result in poor coverage and reduced grass control requiring repeat applications. Spray pressures should reflect a minimum of 30 psi and a maximum of 60 psi at the nozzle. Do not use flood nozzles.

Applications to garlic or onions (dry bulb and green) should be made in a minimum of 20 gals of spray solution per acre.

Air Application

Use a minimum of 3 gals of spray solution per acre unless otherwise directed in this label. Increase spray volumes up to 10 gals as grass or crop foliage becomes dense. For garlic or onions (dry bulb and green): When applying by air do not exceed 16 fl oz/A in a single application. In California, air applications to garlic or onion should be made in a minimum of 20 gals of spray solution per acre. In states other than California, air application to garlic or onion should be made in a minimum of 10 gals of spray solution.

NOTE: Crop injury may occur when V-101371 EC Herbicide is applied to garlic or onion with aerial equipment.

Spot Treatment

When using hand sprayers or high volume sprayers utilizing hand guns, mix 1/3 to 2/3% (0.44 oz to 0.85 oz per gal.) V-101371 EC Herbicide and treat to wet vegetation, while not allowing runoff of spray solution. For uses requiring crop oil concentrate, include crop oil concentrate at 1% (1.3 oz per gal) by volume. For uses requiring non-ionic surfactant, include non-ionic surfactant at 1/4% (0.33 oz per gal) by volume.

NOTE: If V-101371 EC Herbicide is applied as a spot treatment care should be taken to not exceed the maximum rate allowed on a "per acre" basis or crop injury may occur.

CHEMIGATION – GARLIC, ONIONS (Dry Bulb) and GREEN ONION SPRINKLER IRRIGATION APPLICATION

May be applied to onions and garlic by sprinkler irrigation systems.

Apply V-101371 EC Herbicide at the high rate prescribed for annual grasses (32 fl. oz/A) when the grass height is at the high end of the range (application to larger grasses may not provide adequate control). Add a crop oil concentrate containing at least 15% emulsifier at 1 quart per acre or non-ionic surfactant with at least 80% active ingredient at 0.25% v/v of total spray solution.

Apply V-101371 EC Herbicide in 0.1 to 0.2 acre inch of water either at the end of a regular irrigation set or as a separate application not associated with a regular irrigation using the least amount of water that provides proper distribution and coverage. Application of more than label recommended quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness. Use a metering device to inject the V-101371 EC Herbicide into the irrigation water at a constant flow. Constant agitation must be maintained in the chemical supply tank during the entire period of herbicide application. Inject the product with a positive displacement pump into the main line ahead of a right angle turn to ensure adequate mixing. Allow time for all lines to flush the herbicide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of remaining herbicide, a dye indicator may be injected into the lines to mark the end of the application period.

It is not recommended that V-101371 EC Herbicide be applied through an irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Restrictions

- **Do not apply V-101371 EC Herbicide by chemigation in the states of Idaho, Montana, Oregon and Washington.**
- Do not apply by chemigation to any other crop, or to this crop using any other type of irrigation system.
- Do not apply this product through any other type of irrigation system.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

Use Precautions

1. Apply this product only through irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, travelers, big gun, solid set or hand move.
2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.
3. If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
4. A person knowledgeable of chemigation system and responsible for its operation or under supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
5. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
6. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
7. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
8. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
9. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
10. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

SPRAY DRIFT MANAGEMENT

Restrictions

- Do not allow spray from ground or aerial equipment to drift onto adjacent land or crops. When drift may be a problem, do everything possible to reduce spray drift, including:
- Do not apply when conditions are favorable for drift (high temperatures, drought and low relative humidity), especially when sensitive plants are located nearby.
- Do not spray if wind speed is 10 mph or greater. If sensitive crops or plants are downwind, extreme caution must be used under all conditions.
- Do not spray if winds are gusty.
- Do not apply when a temperature inversion exists. If inversion conditions are suspected, consult with local weather services before making an application.
- Do not allow V-101371 EC Herbicide to come in contact with desirable grass crops such as corn, rice, small grains, sorghum, or turf, as these and other grass crops will be injured or killed.
- Do not use nozzles that produce small droplets.
- **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

Further reductions in drift can be obtained by:

1. Using large droplet size sprays. Orient nozzles downward and slightly backward as needed to reduce drift for ground applications.
2. Orienting nozzles straight back with the windstream, using straight stream orifices for aerial applications. Use the lowest number of nozzles practical with the largest possible orifice size to obtain the minimum 3 GPA volume. Application height and boom length should be set according to manufacturer's instructions to minimize drift.
3. Increasing the volume of spray mixture (for example a minimum of 10 GPA for ground applications) by using higher flow rate nozzles. Using lower pressure with the appropriate nozzle to obtain higher volumes will also reduce drift.
4. Applying as close to target plants as practical while maintaining a good spray pattern for adequate coverage.

RESISTANCE MANAGEMENT

V-101371 EC Herbicide is a Group 1 herbicide. Any weed population may contain or develop plants naturally resistant to *V-101371* EC Herbicide and other Group 1 herbicides. Weed species with acquired resistance to Group 1 may eventually dominate the weed population if Group 1 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by *V-101371* EC Herbicide or other Group 1 herbicides. Repeated use of *V-101371* EC Herbicide (or similar postemergence grass herbicide with the same mode of action) may lead to the selection of naturally occurring biotypes that are resistant to these products in some grass species.

If poor performance occurs and cannot be attributed to adverse weather or application conditions, a resistant biotype may be present. This is most likely to occur in fields where other control strategies such as crop rotation, mechanical removal and other classes of herbicides are not used from year to year.

To delay herbicide resistance consider:

- Avoiding the consecutive use of *V-101371* EC Herbicide or other target site of action Group 1 herbicides that have similar target site of action, on the same weed species.
- Using tank mixtures or premixes with herbicides from different target site of action groups as long as the involved products are all registered for the same use, have different sites of action, and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Basing herbicide use on a comprehensive IPM program.
- Monitoring treated weed populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
Alfalfa, Seedling	15 days before grazing, feeding or harvesting (cutting) for forage or hay	9 to 16 fl oz	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v or Crop Oil Concentrate/Methylated Seed Oil at 1 qt/A or 1% v/v See tank mix label for specific adjuvant recommendations.	2.5 to 4 lb/A	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.
Alfalfa, Established including: Sainfoin, Holy clover, Birdsfoot trefoil	15 days before grazing, feeding or harvesting (cutting) for forage or hay	12 to 16 fl oz	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v or Crop Oil Concentrate/Methylated Seed Oil at 1 qt/A or 1% v/v See tank mix label for specific adjuvant recommendations.	2.5 to 4 lb/A	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.
Artichoke (Globe)	5 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.
Asparagus	1 day	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.

continued

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
<p>[Dried Shelled Pea and Bean (except Soybean) Subgroup 6C] Bean, Dry except soybean including: Bean (<i>Lupinus</i> spp.) Grain Sweet White White Sweet Bean (<i>Phaseolus</i> spp.) Field Kidney Lima (dry) Navy Pinto Tepary Bean (<i>Vigna</i> spp.) Adzuki Bean Black-eyed Pea Catjang Cowpea Crowder Pea Moth Bean Mung Bean Rice Bean Southern Pea Urd Bean Broad (dry) Chickpea (garbanzo) Guar Lablab Bean Lentil</p>	30 days	9 to 16 fl oz ⁽⁵⁾	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) peryear.
<p>[Succulent Shelled Pea and Bean Subgroup 6B] Bean, Succulent Shelled including: Bean (<i>Phaseolus</i> spp.) Broad Bean (succulent) Lima Bean (green) Bean (<i>Vigna</i> spp.) Black-eyed Pea Cowpea Southern Pea</p>	21 days	9 to 16 fl oz ⁽⁵⁾	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not apply more than one (1) application per acre per year.
<p>Beet, Garden</p>	30 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) peryear.

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
<p>Berry Low Growing (except Cranberry and Strawberry) [Subgroup 13-07G] including: Bearberry Bilberry Blueberry, lowbush Cloudberry Lingonberry Muntries Partridgeberry</p>	45 days	9 to 16 fl oz ⁽⁵⁾	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>For repeat applications make a minimum of a 14 day interval.</p> <p>Verify Crop tolerance to V-101371 EC Herbicide on a small area of the crop, at the desired V-101371 EC Herbicide rate and with the same Non-ionic Surfactant (NIS) that will be used on the field. If no crop response is evident seven (7) days after treatment, V-101371 EC Herbicide may be used on the entire field at the rate tested and with the same NIS used in the tolerance test.</p> <p>Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application.</p> <p>Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.</p>
<p>[Head & Stem Brassica Subgroup 5A] Brassica Vegetables, Head and Stem including: Broccoli Brussels Sprouts Cabbage Cauliflower Cavalo Broccolo Chinese broccoli; Chinese (Napa) Cabbage Chinese Mustard Kohlrabi.</p>	30 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>For repeat applications make on a minimum of a 14 day interval</p> <p>Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application.</p> <p>Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.</p>

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
<p>[Bushberry Subgroup 13-07B] Bushberry including: Aronia Berry Blueberry, Highbush Chilean Guava Cranberry, Highbush Currant, Black Currant, Buffalo Currant, Native Currant, Red Elderberry European Barberry Gooseberry Honeysuckle Edible Huckleberry Jostaberry Juneberry, Saskatoon Berry Native Currant Salal Sea Buckthorn Cultivars, varieties and/or hybrids of these.</p>	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.
<p>[Caneberry Subgroup 13-07A] Caneberry including: Blackberry Loganberry Raspberry, Black Raspberry, Red Raspberry, Wild Cultivars, varieties and/or hybrids of these.</p>	7 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
<p>Canola [including Rapeseed Subgroup 20A, except flax seed, mustard seed and sesame seed]</p> <p>(Not for use in California)</p> <p>Borage Crambe Cuphea Echium Gold of Pleasure (Camelina) Hare's Ear Mustard Lesquerella Lunaria Meadowfoam Milkweed Oil Radish Poppy seed Rapeseed (canola) Sweet Rocket</p>	70 days	9 to 12 fl oz ⁽⁵⁾	12 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>Verify Crop tolerance to V-101371 EC Herbicide on a small area of the crop, at the desired V-101371 EC Herbicide rate and with the same Non-ionic Surfactant (NIS) that will be used on the field. If no crop response is evident seven (7) days after treatment, V-101371 EC Herbicide may be used on the entire field at the rate tested and with the same NIS used in the tolerance test.</p> <p>Restrictions Do not apply more than 12 fl oz/A (0.091 lb ai/A) per application.</p> <p>Do not apply more than 12 fl oz/A (0.091 lb ai/A) per year.</p> <p>Do not apply after crop has begun bolting. Crop injury may occur when V-101371 EC Herbicide is applied during the bloom period</p>
Carrot	30 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>For repeat applications make on a minimum of a 14 day interval.</p> <p>Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application.</p> <p>Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.</p>

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
Clover	15 days before grazing, feeding or harvesting (cutting) for forage or hay	9 to 16 fl oz	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	2.5 to 4 lb/A	For use on clover grown in the states of Idaho, Oregon and Washington only. For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) per application. Do not apply more than 32 fl oz/A(0.242 lb ai/A) per year.
Corn, Field ⁽⁶⁾	90 days	6 fl oz	-	Non-ionic surfactant (NIS) at 0.25% v/v plus Ammonium Sulfate Do not use Crop Oil Concentrate or Methylated Seed Oil with V-101371 EC Herbicide in this use pattern.	2.5 to 4 lb/A	To control existing stand, replant no sooner than 6 days after application. Restrictions Do not make more than 1 application per year. Do not apply more than 6 fl oz/A (0.045 lb ai/A) per year.
Cotton (including cotton grown for seed)	60 days	9 to 16 fl oz	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v or Crop Oil Concentrate/Methylated Seed Oil at 1 qt/A or 1% v/v See tank mix label for specific adjuvant recommendations	2.5 to 4 lb/A	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year. Do not graze treated fields or feed treated forage or hay to livestock.
Cranberry	30 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year. Do not apply between the "hook" stage and full fruit set.

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
[Cucurbit Vegetables Crop Group 9] Cucurbits including: Cantaloupes (all) Cucumber Gherkin Honeydew Melon Muskmelons (all) Pumpkin Squash (all) Watermelon	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.
Fallow Land Conifer Trees (and other non-producing agricultural areas) Non-Crop or Non-Planted Areas	N/A	9 to 16 fl oz	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v or /Crop Oil Concentrate/ Methylated Seed Oil at 1 qt/A or 1% v/v	2.5 to 4 lbs/A	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) per application. Do not apply more than 64 fl. oz/A (0.485 lb ai/A) per year. Do not plant any crop for 30 days after application unless clethodim is registered for use in that crop.
Flax (Not for use in California)	60 days	9 to 16 fl oz ⁽⁵⁾	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	Apply prior to bloom. Crop injury may occur when V-101371 EC Herbicide is applied during the bloom period. For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not apply more than 32 fl oz (0.242 lb ai/A) per year.

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
<p>[Fruiting Vegetables (except Tomato) Crop Group 8-10] Fruiting Vegetable (except Tomato) including: African Eggplant Bush Tomato Bell Pepper Cocona Currant Tomato Eggplant Garden Huckleberry Goji Berry Groundcherry Martynia Naranjilla Okra Pea Eggplant Pepino Nonbell Pepper; Roselle Scarlet Eggplant Sunberry Tomatillo Tree Tomato cultivars, varieties, and/or hybrids of these.</p>	20 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>For repeat applications make on a minimum of a 14 day interval.</p> <p>Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application.</p> <p>Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.</p>
<p>[Subgroup 19A] Herbs including: Angelica Balm Basil Borage Burnet Camomile Catnip Chervil (dried) Chive Chive, Chinese Clary Coriander (leaf) Costmary Culantro (leaf) Curry (leaf) Dill (dillweed) Horehound Hyssop Lavender Lovage (leaf) Marigold Marjoram (<i>Origanum</i> spp.) Nasturtium Parsley (dried) Pennyroyal Rosemary Rue Sage Savory, Summer and Winter Sweet Bay Tansy Tarragon Thyme Wintergreen Woodruff Wormwood</p>	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>For repeat applications make on a minimum of a 14 day interval.</p> <p>V-101371 EC Herbicide has not been tested on all herbs, and herb varieties. It is the responsibility of the user to test V-101371 EC Herbicide on a small portion of the crop to be treated before treating the entire field.</p> <p>Verify crop tolerance to V-101371 EC Herbicide on a small area of the crop, at the desired V-101371 EC Herbicide rate and with the same crop oil concentrate that will be used on the field. If no crop response is evident seven (7) days after treatment, V-101371 EC Herbicide may be used on the entire field at the rate tested and with the same crop oil used in the tolerance test.</p> <p>Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application.</p> <p>Do not apply more than 64 fl oz/A (0.485 lb. ai/A) per year.</p>

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
Hops	21 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat application make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.
Leaf Lettuce	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.
[Subgroup 4B] Leaf Petioles including: Cardoon Celery Celtuce Chinese Celery Fennel, Florence (finocchio) Rhubarb Swiss Chard	30 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.
[Subgroup 5B] Leafy Brassica Greens including: Broccoli Raab Cabbage, Chinese (bok choy) Collards Kale Mizuna Mustard Greens Mustard Spinach Rape Greens	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat application make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
<p>[Subgroup 4A] Leafy Greens including: Amaranth Chinese Spinach Leafy Amaranth Tampala Arugula (roquette) Chervil Chrysanthemum, Edible-leaved Chrysanthemum, Garland Corn Salad Cress Garden Upland (yellow rock and winter) Dandelion Dock (sorrel) Endive (escarole) Lettuce, Head and Leaf Orach Parsley Purslane Garden Winter Radicchio (red chicory) Spinach Spinach New Zealand Vine (Indian and Malabar)</p>	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>For repeat applications make on a minimum of a 14 day interval.</p> <p>Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) application.</p> <p>Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.</p>
<p>[Subgroup 6A] Legume Vegetables, Edible Podded including: Bean (<i>Phaseolus</i> spp.) Runner Snap Wax Bean (<i>Vigna</i> spp.) Asparagus Chinese Longbean Moth Yardlong Jackbean Edamame, immature soybean Pea (<i>Pisum</i> spp.) Dwarf Edible-pod Snow Sugar Snap Pigeon Sword Bean</p>	21 days	9 to 16 fl oz ⁽⁵⁾	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>For peas apply before bloom, but no later than 21 days before harvest.</p> <p>Restrictions Do not apply more than 16 fl oz/A(0.121 lb ai/A) per application.</p> <p>Do not apply more than one (1) application per acre per year.</p>

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
Mint	21 days	9 to 16 fl oz	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v or Crop Oil Concentrate/Methylated Seed Oil at 1 qt/A or 1%v/v	2.5 to 4 lb/A	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.
Mustard Seed (Not for use in California)	75 days	9 to 12 fl oz ⁽⁵⁾	12 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply after crop has begun bolting. Crop injury may occur when V-101371 EC Herbicide is applied during the bloom period. Do not apply more than 12 fl oz/A (0.091 lb ai/A) per year.
[Bulb Onion Subgroup 3-07A] Onions (Dry Bulbs Only) ^{7, 8)} Daylily, Bulb Fritillaria Bulb Garlic, Bulb Garlic, Great-headed, Bulb Garlic, Serpent, Bulb Lily Bulb Onion, Bulb Onion, Chinese, Bulb Onion, Pearl Onion, Potato, Bulb Shallot, Bulb cultivars, varieties, and/or hybrids of these.	45 days	9 to 16 fl oz	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Minimum of 20 gals/A spray volume by ground in entire U.S. Minimum of 20 gals/A spray volume by air in California. In states other than California, air applications to onions or garlic should be made in a minimum of 10 gals/A. Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
Onions, Green ^{7,8)}	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>For repeat applications make on a minimum of a 14 day interval.</p> <p>Minimum of 20 gals/A spray volume by air in California⁽⁵⁾</p> <p>In states other than California, air applications to onions or garlic should be made in a minimum of 10 gals/A.⁽⁷⁾</p> <p>Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) in a single application.</p> <p>Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.</p>
Ornamentals	N/A	9 to 16 fl oz	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>For repeat applications make on a minimum of a 14 day interval.</p>
Non-Bearing Food Crops	N/A	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>Sugar maples cannot be tapped for syrup within one year of V-101371 EC Herbicide application.</p> <p>Restrictions Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.</p> <p>Do not apply more than 16 fl oz/A (0.121 lb ai/A) in a single application to non-bearing food crops.</p>
[Subgroup 6C] Pea, Shelled (<i>Pisum</i> spp.) Field Pigeon	30 days	9 to 16 fl oz ⁽⁵⁾	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>Apply before bloom but not later than 30 days prior to harvest.⁽⁹⁾</p> <p>Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application.</p> <p>Do not apply more than one (1) application per acre per year.</p>

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
[Subgroup 6B] Pea, Succulent Shelled (<i>Pisum</i> spp.) English Pea Garden Pea Green Pea Pigeon Pea	21 days	9 to 16 fl oz ⁽⁵⁾	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	Apply before bloom but not later than 21 days prior to harvest. ⁽⁹⁾ Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not apply more than one (1) application per acre per year.
Peanut (including perennial)	40 days	9 to 16 fl oz	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v or Crop Oil Concentrate/Methylated Seed Oil at 1 qt/A or 1% v/v	2.5 to 4 lbs/A	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) in a single application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year, except in Florida where the limit is 64 fl oz/A (0.485 lb ai/A) per year (perennial peanuts only).
[Crop Group 11-10] Pome Fruit (including) Apple Azarole Crabapple Loquat Mayhaw Medlar Pear Pear, Asian Quince Quince, Chinese Quince, Japanese Tejocote	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) in a single application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.
Potato	30 days	9 to 16 fl oz	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v or Crop Oil Concentrate/Methylated Seed Oil at 1 qt/A or 1% v/v	2.5 to 4 lbs/A	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) in a single application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.
Radish	15 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) in a single application. Do not apply more than 32 fl oz/A (0.242 lb ai/A)

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
						per year.
[Subgroup 1A] Root Vegetables (except Sugar Beet and Radish) ⁽¹⁰⁾	30 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) in a single application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
Safflower	70 days	9 to 16 fl oz	12-16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>For repeat applications make on a minimum of a 14 day interval.</p> <p>Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) in a single application.</p> <p>Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.</p>
Sesame	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>For repeat applications make on a minimum of a 14 day interval.</p> <p>Restrictions Do not apply during flowering.</p> <p>Do not apply more than 16 fl oz/A (0.121 lb ai/A) in a single application.</p> <p>Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.</p>
Soybean ⁽¹¹⁾	60 days	9 to 16 fl oz ⁽⁵⁾	12 to 32 fl oz	<p>Non-ionic Surfactant (NIS) at 0.25% v/v or Crop Oil Concentrate/ Methylated Seed Oil at 1 qt/A or 1% v/v</p> <p>See tank mix label for specific adjuvant recommendations.</p>	2.5 to 4 lbs/A	<p>For repeat applications make on a minimum of a 14 day interval.</p> <p>Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) per application.</p> <p>Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.</p> <p>Do not graze treated fields or feed treated forage or hay to livestock.</p>

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-10137 1 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
Stevia (dried leaves)	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>For repeat applications make on a minimum of a 14 day interval.</p> <p><i>V-10137 1 EC</i> Herbicide has not been tested on all varieties. It is the responsibility of the user to test <i>V-10137 1 EC</i> Herbicide on a small portion of the crop to be treated before treating the entire field.</p> <p>Verify crop tolerance to <i>V-10137 1 EC</i> Herbicide on a small area of the crop, at the desired <i>V-10137 1 EC</i> Herbicide rate and with the same Non-ionic Surfactant (NIS) that will be used on the field. If no crop response is evident seven (7) days after treatment, <i>V-10137 1 EC</i> Herbicide may be used on the entire field at the rate tested and with the same NIS used in the tolerance test.</p> <p>Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) in a single application.</p> <p>Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.</p>

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
<p>[Crop Group 12-12] Stone Fruit including Apricot Apricot, Japanese Capulin Cherry, Black Cherry, Nanking Cherry, Sweet Cherry, Tart; Jujube, Chinese Nectarine Peach Plum Plum, American Plum, Beach Plum, Canada Plum, Cherry Plum, Chickasaw Plum, Damson Plum, Japanese; Plum, Klamath Plum, Prune Plumcot Sloe</p>	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) in a single application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.
<p>Strawberry</p>	4 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) in a single application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
Sugar Beet	40 days	9 to 16 fl oz ⁽⁵⁾	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v or Crop Oil Concentrate/ Methylated Seed Oil at 1 qt/A or 1% v/v See tank mix label for specific adjuvant recommendations.	2.5 to 4 lbs/A	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.
[Subgroup 20B] Sunflower including: Calendula Casteroil Plant Chinese Tallowtree Euphorbia Evening Primrose Jojoba Niger Seed Rose Hip Stokes Aster Tallowwood Tea Oil Plant Vernonia	70 days	9 to 16 fl oz ⁽⁵⁾	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v or Crop Oil Concentrate/ Methylated Seed Oil at 1 qt/A or 1% v/v See tank mix label for specific adjuvant recommendations.	2.5 to 4 lbs/A	For repeat applications make on a minimum of a 14 day interval. Verify crop tolerance to V-101371 EC Herbicide on a small area of the crop, at the desired V-101371 EC Herbicide rate and with the same crop oil concentrate that will be used on the field. If no crop response is evident seven (7) days after treatment, V-101371 EC Herbicide may be used on the entire field at the rate tested and with the same crop oil used in the tolerance test. Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.
[Subgroup 1D] Sweet Potato, Yam and other tuberous and corm vegetables (except potato) ⁽¹²⁾	30 days	9 to 16 fl oz	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.
Tomato	20 days	9 to 16 fl oz	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.

Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-101371 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
Turnip Greens	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat application make on a minimum of a 14 day interval. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) in a single application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.

N/A = Not Applicable

(1) V-101371 EC Herbicide is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided.

(2) See annual and perennial grass control tables for specific use rate recommendations.

(3) Non-ionic surfactant (NIS) in this case refers to an adjuvant containing at least 80% non-ionic surfactant. Crop oil concentrate in this case refers to both crop oil concentrate and crop oil concentrate blends. **Acceptable crop oil concentrates** would be those that contain a minimum of 80% oils and 15% emulsifier. **Acceptable crop oil concentrate blends** would be those that contain a minimum of 60% oils and 25-40% surfactants and emulsifiers. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils.

(4) Use spray grade ammonium sulfate. The use of ammonium sulfate does not take the place of the required adjuvant.

(5) See DIRECTIONS FOR REDUCED RATE USE TO CONTROL SMALL ANNUAL GRASSES table.

(6) For burndown of existing stand of Roundup Ready field corn or volunteer Roundup Ready field corn prior to replanting field corn. See RECOMMENDATIONS FOR USE IN ROUNDUP READY FIELD CORN (BURNDOWN) table.

(7) If V-101371 EC Herbicide is applied as a spot treatment to garlic, onion or non-bearing food crops care should be taken to not exceed the maximum rate allowed on a "per acre" basis or crop injury may occur.

(8) In California, do not apply V-101371 EC Herbicide to garlic or onions until crop has at least two full leaves. In California, 14 days spray intervals are recommended between the application of V-101371 EC Herbicide and liquid nitrogen or other herbicide applications. Injury to crop may occur when shorter intervals are observed.

(9) Applications of V-101371 EC Herbicide to peas during the bloom period could result in severe crop injury, including loss of yield and delayed maturity.

(10) Other root vegetables approved for use with V-101371 EC Herbicide include: burdock, edible; celeriac; chervil, turnip-rooted; chicory; ginseng; parsley, turnip-rooted; parsnip; radish, oriental; rutabaga; salsify; salsify, black; salsify, Spanish; skirret and turnip.

(11) See V-101371 EC HERBICIDE TANK MIX WITH BROADLEAF HERBICIDES FOR THE CONTROL OF VOLUNTEER CORN (INCLUDING ROUNDUP READY) IN SOYBEAN.

(12) Other tubereous and corm vegetables approved for use with V-101371 EC Herbicide include: arracacha, arrowroot, Chinese artichoke, Jerusalem artichoke, edible canna, cassava, bitter and sweet chayote (root), chufa, dasheen (taro), ginger, leren, tanier, turmeric and bean yam.

USE DIRECTIONS FOR ANNUAL GRASSES ALL CROPS

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

Restrictions

- Do not exceed the maximum per application rate listed in Table 1, CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-10137 1 EC HERBICIDE.
- Do not exceed the maximum yearly rate listed in Table 1, CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-10137 1 EC HERBICIDE.

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (inches)	APPLICATION RATES	
			MINIMUM RATE fl oz/A	MAXIMUM RATE ⁽¹⁾ fl oz/A
Barnyardgrass	<i>Echinochloa crus-galli</i>	2 to 8	9	16
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>	2 to 6	9	16
Brome				
California	<i>Bromus carinatus</i>	2 to 6	9	16
Cheat	<i>Bromus secalinus</i>	2 to 6	9	16
Downy	<i>Bromus tectorum</i>	2 to 6	9	16
Ripgut	<i>Bromus diandrus</i>	2 to 6	9	16
Canarygrass	<i>Phalaris canariensis</i>	1 to 4	9	16
Crabgrass				
Hairy	<i>Digitaria adscendens</i>	2 to 6**	9	16
Large	<i>Digitaria sanguinalis</i>	2 to 6**	9	16
Smooth	<i>Digitaria ischaemum</i>	2 to 6**	9	16
Southern	<i>Digitaria ciliaris</i>	2 to 6**	9	16
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	2 to 6**	9	16
Fall Panicum	<i>Panicum dichotomiflor</i>	2 to 8	9	16
Field Sandbur	<i>Cenchrus incertus</i>	2 to 6	9	16
Foxtail				
Giant	<i>Setaria faberi</i>	2 to 12	9	16
Green	<i>Setaria viridis</i>	2 to 8	9	16
Yellow	<i>Setaria glauca</i>	2 to 8	9	16
Goosegrass	<i>Eleusine indica</i>	2 to 6**	9	16
Itchgrass	<i>Rottboellia cochinchinensis</i>	2 to 6	9	16
Junglerice	<i>Echinochloa colona</i>	2 to 6	9	16
Lovegrass (Stinkgrass)	<i>Eragrostis cilianensis</i>	2 to 6	9	16
Rabbitsfootgrass	<i>Polypogon monspeliensis</i>	1 to 4	9	16
Red Rice	<i>Oryza sativa</i>	1 to 3	9	16
Rygrass				
Hardy	<i>Lolium remotum</i>	2 to 6	9	16
Italian	<i>Lolium multiflorum</i>	2 to 6	9	16
Seedling Johnsongrass	<i>Sorghum halepense</i>	4 to 10	9	16
Shattercane	<i>Sorghum bicolor</i>	6 to 18	9	16
Southwestern Cupgrass	<i>Eriochloa gracilis</i>	2 to 6	9	16
Sprangletop				
Amazon	<i>Leptochloa panicoides</i>	2 to 6	9	16
Bearded	<i>Leptochloa fascicularis</i>	2 to 6	9	16
Mexican	<i>Leptochloa uninervia</i>	2 to 6	9	16
Red	<i>Leptochloa filiformis</i>	2 to 6	9	16

continued

USE DIRECTIONS FOR ANNUAL GRASSES
ALL CROPS (continued)

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (inches)	APPLICATION RATES	
			MINIMUM RATE fl oz/A	MAXIMUM RATE ⁽¹⁾ fl oz/A
Texas Panicum Volunteer Cereals ⁽³⁾	<i>Panicum texanum</i>	2 to 6	9	16
Barley	<i>Hordeum vulgare</i>	2 to 6	9	16
Oats	<i>Avena sativa</i>	2 to 6	9	16
Rye	<i>Secale cereale</i>	2 to 6	9	16
Wheat ⁽²⁾	<i>Triticum aestivum</i>	2 to 6	9 ⁽²⁾	16
Volunteer Corn ^(2,3)	<i>Zea mays</i>	up to 12	6	12
Volunteer Corn ⁽³⁾	<i>Zea mays</i>	up to 24	9	14
Volunteer Corn ^(2,3)	<i>Zea mays</i>	up to 36	12	16
Volunteer Grain Sorghum	<i>Sorghum bicolor</i>	8 to 12	9	16
Wild Oats	<i>Avena fatua</i>	2 to 6	9	16
Wild Proso Millet	<i>Panicum miliaceum</i>	2 to 10	9	16
Witchgrass	<i>Panicum capillare</i>	2 to 8	9	16
Woolly Cupgrass	<i>Eriochloa villosa</i>	2 to 8	9	16

*Generally occurs between 3-leaf stage and tillering.

**Length of lateral growth.

- (1) Rates higher than 16 fl oz/A may be applied in certain geographic areas, cropping situations or environmental conditions, where experience has shown that higher rates are needed for satisfactory control of annual grasses. In these situations, rates from 16 to 32 fl oz/A may be applied.

Restrictions

Do not apply more than 16 fl oz/A of V-101371 EC Herbicide per application to the following crops: garden beets, berry low growing (except cranberry and strawberry), bushberry, broccoli, cabbage, caneberry, carrot, cauliflower (and other head and stem brassica vegetables), celery, cranberry, cucurbits, flax, fruiting vegetables (except tomato), green onion, herbs, hops, leaf lettuce, legume vegetables, pea succulent shelled, pome fruit, radish (and other root vegetables), rhubarb (and other leaf petioles), safflower, sesame, stevia (dried leaves), stone fruit, strawberry, turnip greens and non-bearing food crops.

Do not apply more than 12 fl oz/A of V-101371 EC Herbicide per application to canola or mustard seed.

- (2) When a cereal grain crop (such as wheat) is interseeded for crop establishment or is planted as wind breaks to aid crop establishment, the minimum V-101371 EC Herbicide use rate for control is 12 fl oz/A.
- (3) Includes Roundup Ready, Liberty Link® and Clearfield® volunteer corn; however not Sethoxydim-Resistant volunteer corn.

**USE DIRECTIONS FOR ANNUAL & PERENNIAL GRASS CONTROL
IN ESTABLISHED ALFALFA AND MINT WITH V-10137 1 EC HERBICIDE**

GRASS SPECIES	WEED SPECIES AND SIZE	APPLICATION RATES
Annual & Perennial Grasses Listed in Grass Tables	See Tables	See Table 1, CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITAITONS FOR V-10137 1 EC HERBICIDE

Mowing: The best control of annual grasses can be achieved by applying V-10137 1 EC Herbicide before grass weeds are mowed. Once a grass is mowed it becomes tougher to control, as much of the available leaf surface has been removed. In areas without a killing frost, some annuals can over-winter after having been mowed multiple times. These grasses form large crowns and may contain many viable buds. These grasses, even though they may be an annual grass, may require repeated applications of V-10137 1 EC Herbicide for partial or complete control.

Irrigated Alfalfa and Mint: Irrigation practices can be very critical to the successful use of V-10137 1 EC Herbicide in established alfalfa and mint and may be necessary to initiate active growth of the weeds prior to application. Generally applications 2 to 4 days after an irrigation are most effective. Irrigation made shortly after application (2 days) can be effective, but more consistent grass control occurs when the irrigation is made before the application.

Aerial Application: Apply V-10137 1 EC Herbicide in a minimum of 10 GPA in established alfalfa and mint when applying by air.

Annual Grass Control: Apply V-10137 1 EC Herbicide at the grass sizes indicated in the Recommendation for Annual Grass Table and rates indicated. If a grass has been cut, apply V-10137 1 EC Herbicide after active growth has resumed and regrowth has reached the minimum height and before it reaches the maximum height indicated. Apply before the alfalfa/mint canopy covers the grasses and interferes with the spray coverage. Some annual grasses are spring- and summer- germinating plants, while others are fall-germinating plants, and the time they are actively growing and most susceptible to V-10137 1 EC Herbicide may vary from region to region. Also some annuals germinate over an extended period of time, and because control of small grasses is desired, applications after each weed flush may be required. As a general rule spray spring and summer germinating grasses as early in the season as possible, after initial green-up. Spray fall -germinating weeds in the fall soon after they begin growing but before any damage is done due to frost. Late fall applications may be less effective due to environmental conditions, such as frost, slower plant growth or the onset of flowering.

Perennial Grass Control: V-10137 1 EC Herbicide effectively controls perennial grasses such as bermudagrass, Johnsongrass, quackgrass, wirestem muhly, tall fescue, foxtail barley and orchardgrass. Due in part to lack of tillage, perennial grasses are more difficult to control in a perennial crop such as established alfalfa or mint. A program of repeated applications is usually necessary for best results. The best way to control perennial grasses is to do so in the year of stand establishment before rhizomes and stolons become large and difficult to kill.

Use the high rate under heavy grass pressure and/or when grasses are at or near maximum height.

**DIRECTIONS FOR REDUCED RATE USE TO CONTROL SMALL ANNUAL GRASSES
CANOLA, FLAX, LEGUME VEGETABLES (DRY AND SUCCULENT), MUSTARD SEED,
SOYBEAN, SUGAR BEET AND SUNFLOWER
(REDUCED RATE DIRECTIONS NOT FOR USE IN CALIFORNIA)**

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Regrowth by tillering may occur if application is made when plants are stressed by lack of moisture, excessive moisture, low or high temperatures and/or under very low humidity.

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT (inches)	RATE FL OZ/ ACRE⁽¹⁾
Barnyardgrass	<i>Echinochloa crus-galli</i>	1 to 4	6
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>	1 to 4	8
Crabgrass			
Large	<i>Digitaria sanguinalis</i>	1 to 3*	6
Large	<i>Digitaria sanguinalis</i>	1 to 4*	8
Smooth	<i>Digitaria ischaemum</i>	1 to 3*	6
Smooth	<i>Digitaria ischaemum</i>	1 to 4*	8
Southern	<i>Digitaria ciliar</i>	1 to 4*	8
Fall Panicum	<i>Panicum dichotomiflorum</i>	1 to 4	6
Foxtail			
Giant	<i>Setaria faberi</i>	1 to 4	6
Green	<i>Setaria viridis</i>	1 to 4	6
Millet	<i>Setaria italica</i>	1 to 4	8
Yellow	<i>Setaria glauca</i>	1 to 4	6
Seedling Johnsongrass	<i>Sorghum halepense</i>	1 to 6	8
Shattercane	<i>Sorghum bicolor</i>	4 to 10	6
Texas Panicum	<i>Panicum texanum</i>	1 to 4	8
Volunteer Cereals			
Barley	<i>Hordeum vulgare</i>	1 to 4	8
Oats	<i>Avena sativa</i>	1 to 4	8
Wheat	<i>Triticum aestivum</i>	1 to 4	8
Volunteer Corn**	<i>Zea mays</i>	4 to 12	6
Wild Proso Millet	<i>Panicum miliaceum</i>	1 to 6	6
Wild Oats	<i>Avena fatua</i>	1 to 4	8

*Length of lateral growth

**Not S.R. Corn

⁽¹⁾Always add a non-ionic surfactant at 0.25% v/v total spray volume unless crop specific restrictions and limitations advise otherwise.

**USE DIRECTIONS FOR PERENNIAL GRASSES
(ALL CROPS)**

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

Restrictions

- Do not exceed the maximum per application rate listed in Table 1, CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-10137 1 EC Herbicide.
- Do not exceed the maximum yearly rate listed in Table 1, CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-10137 1 EC HERBICIDE.

GRASS SPECIES	WEED HEIGHT (inches)	APPLICATION RATE	
		MINIMUM RATE fl oz/A	MAXIMUM RATE fl oz/A
Bermudagrass (<i>Cynodon dactylon</i>) First Application	3 (or up to 6" runners)	12	32
Repeat Application(s) (if regrowth occurs)	3 (or up to 6" runners)	12	32
Fescue, Tall (<i>Festuca arundinacea</i>) First Application	4 to 8	12	32
Repeat Application(s) (if regrowth occurs)	4 to 8	12	32
Foxtail Barley (<i>Hordeum jubatum</i>) First Application	2 to 6	12	32
Repeat Application (if regrowth occurs)	2 to 6	12	32
Orchardgrass (<i>Dactylis glomerata</i>) First Application	4 to 8	12	32
Repeat Application(s) (if regrowth occurs)	4 to 8	12	32
Quackgrass (<i>Elytrigia repens</i>) First Application	4 to 12	12	32
Repeat Application(s) (if regrowth occurs)	4 to 12	12	32
Rhizome Johnsongrass (<i>Sorghum halepense</i>) First Application	12 to 24	12	32
Repeat Application(s) (if regrowth occurs)	6 to 18	9	24
Wirestem Muhly (<i>Muhlenbergia frondosa</i>) First Application	4 to 8	12	32
Repeat Application(s) (if regrowth occurs)	4 to 8	12	32
Perennial Bluegrass [Roughstalk (<i>Poa trivialis</i>)] [Kentucky (<i>Poa prantensis</i>)] First Application	2 to 4	12	32
Repeat Application(s)	2 to 4	12	32
Bentgrass (<i>Agrostis</i> spp.) First Application	2 to 4	-	32
Repeat Application(s) (if regrowth occurs)	2 to 4	-	32

**USE DIRECTIONS FOR ANNUAL BLUEGRASS CONTROL
WITH V-10137 1 EC HERBICIDE
ALL CROPS**

GRASS SPECIES	WEED STAGE	APPLICATION RATES	
		MINIMUM RATE fl oz/A	MAXIMUM RATE fl oz/A
Annual Bluegrass (Poa annua)	to 4-leaf	12*	**

Apply under favorable soil moisture and humidity, which exists within a few days after rainfall or within 7 days after irrigation. Grass needs to be actively growing at time of application(s).

Apply at weed stage indicated on the label, as reduced control can be expected with more mature annual bluegrass.

Use the high rate under heavy grass pressure and/or when annual bluegrass is more mature.

See Table 1 for crop specific adjuvant recommendations.

*Use a minimum of 17 fl oz/A to control annual bluegrass in seedling and established alfalfa and mint.

**See Special Use Instructions and Restrictions in Table 1, CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-10137 1 EC Herbicide, for maximum application rates.

USE DIRECTIONS FOR USE IN ROUNDUP READY FIELD CORN (BURNDOWN)

GRASS SPECIES	WEED SIZE (inches)	APPLICATION RATES
		Rate when applied alone or with glyphosate
Field Corn	Up to 12	6 fl oz/A

For control of existing stand of Roundup Ready field corn or volunteer Roundup Ready field corn prior to replanting field corn.

Care must be taken to avoid in field boom (spray) overlaps or excessive crop injury may occur.

Replant no sooner than 6 days after application.

Adjuvant recommendations: Non-ionic Surfactant (NIS) at 0.25% v/v plus AMS at 2.5 to 4 lb/A.

- **Restriction:** Do not use a COC or MSO with V10137 1 EC Herbicide in this use pattern.

TANK MIX - LABEL INFORMATION

The labels for each of the herbicides recommended for tank mixing with *V-10137 1 EC Herbicide* are unique to the characteristics of those products and contain restrictions and limitations that may be more restrictive than the *V-10137 1 EC Herbicide* label in certain considerations. Those concerns may include, but are not limited to:

1. **Always read and follow the restrictions and limitations for all products whether used alone or in a tank mix. The most restrictive labeling of any product used applies in tank mixtures, including all crop rotational and other crop restrictions.**
2. Geographic restrictions - all products are not registered for use in all areas and rates may vary from one region of labeled use to another;
3. Crop rotation restrictions;
4. Applicator certification requirements;
5. Worker safety rules (e.g. protective clothing, reentry time, posting);
6. Soil type or soil characteristics (e.g. pH, OM);
7. Maximum dosage or number of applications per season;
8. Rain free period required; or
9. Application timing (e.g. pre-harvest interval)
10. **Restriction:** Do not exceed the total season rates.

THE MOST RESTRICTIVE LABELING OF ANY PRODUCT USED IN A TANK MIX MUST BE FOLLOWED.

TANK MIX APPLICATION OF *V-10137 1 EC HERBICIDE* AND BROADLEAF HERBICIDES FOR CONTROL OF GRASSES AND BROADLEAF WEEDS

- Apply only to actively growing grass and broadleaf weeds at recommended height or growth stage listed on each label.
- Apply when the first grass or broadleaf weed species in a mixed population reaches the recommended height or growth stage for treatment.
- Apply under favorable soil moisture and humidity that exist a few days after rainfall or within seven days after irrigation.
- Always add the appropriate adjuvant to the spray mix at the rate recommended for each specific tank mix combination.
- Tank mix applications may sometimes result in reduced grass control and possible increases in crop injury as compared to either product used alone. If regrowth occurs, or an additional flush of new grass emerges, make a second application of *V-10137 1 EC Herbicide*, as specified in the respective size and rate tables.
- **Restriction:** Do not tank mix *V-10137 1 EC Herbicide* when broadleaf weeds are tall and/or dense enough to prevent proper grass coverage.

MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, add the correct amount of *V-10137 1 EC Herbicide*. Agitation should create a rippling or rolling action on the water surface.
3. If tank mixing *V-10137 1 EC Herbicide* with other labeled herbicides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions. Prepare no more spray mixture than is required for the immediate spray operation.
4. Add any required adjuvants (crop oil concentrate, non-ionic surfactant and/or nitrogen solution).
5. Fill spray tank to desired level with water. **Agitation should continue until all spray solution has been applied.**

Failure to agitate the spray solution may result in improper mixing of the herbicides and unsatisfactory weed control. Mixing and compatibility qualities should be verified by a jar test.

INFORMATION ON ANTAGONISM

Tank mixes of *V-10137 1 EC Herbicide* with postemergence broadleaf herbicides have shown some reduction or failure to control certain grass species which would have otherwise been controlled when *V-10137 1 EC Herbicide* is applied alone. Activity of the postemergence broadleaf herbicide in the tank mix is not affected.

ALFALFA

**Table 2. V-10137 1 EC HERBICIDE TANK MIXES WITH BROADLEAF HERBICIDES FOR ALFALFA
(Refer to the use directions tables above for specific grasses and growth stages.)**

PRODUCT ⁽¹⁾	APPLICATION RATES/ACRE ⁽²⁾		SPRAY ADDITIVES			
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND APPLICATION		AIR APPLICATION	
			Adjuvant Recommendation	AMS	Adjuvant Recommendation	AMS
V-10137 1 EC Herbicide + 2,4-DB ⁽³⁾	12 to 32 fl oz + Refer to 2,4-DB label	16 to 32 fl oz + Refer to 2,4-DB label	NIS at 0.25% v/v	AMS at 2.5 lb/A	NIS at 0.25% v/v	17 lb/100 gals of spray solution
V-10137 1 EC Herbicide + Pursuit® DG ⁽⁴⁾ or Pursuit ⁽⁴⁾	12 to 32 fl oz + Refer to Pursuit DG or Pursuit labels for use rates	-	NIS at 0.25% v/v	AMS at 2.5 lb/A	NIS at 0.25% v/v	17 lb/100 gals of spray solution
V-10137 1 EC Herbicide + Buctril® 2L ⁽⁵⁾	12 to 32 fl oz + Refer to Buctril 2L label for use rate.	-	NIS at 0.25% v/v	AMS at 2.5 lb/A	NIS at 0.25% v/v	17 lbs/100 gals of spray solution
V-10137 1 EC Herbicide + Raptor®	12 to 32 fl oz + Refer to Raptor label for use rate.	-	NIS at 0.25% v/v	AMS at 2.5 lb/A	NIS at 0.25% v/v	17 lb/100 gals of spray solution

⁽¹⁾ Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.

⁽²⁾ If grass regrowth occurs or an additional flush of new grass emerges, make a second application of V-10137 1 EC Herbicide alone (without a tank mix herbicide), according to the appropriate size and rate prescribed.

⁽³⁾ V-10137 1 EC Herbicide plus 2,4-DB may increase the severity of crop injury when tank mixed. Alfalfa plants will generally outgrow this temporary crop injury within a few weeks.

⁽⁴⁾ Before using this tank mix, read and understand the Pursuit or Pursuit DG labels for geographical restrictions and restrictions regarding alfalfa growth stage and type. Failure to do so can result in crop injury to alfalfa. Do not feed, graze or harvest alfalfa for 30 days following an application of Pursuit to alfalfa.

⁽⁵⁾ In the states of Washington, Oregon, Idaho, Montana, Wyoming, Colorado, Utah, Nevada and the western halves of North Dakota, South Dakota, Nebraska and Kansas: The V-10137 1 EC Herbicide plus Buctril tank mix must be applied in the fall or spring to seedling alfalfa when the majority of the field has a minimum of 2 trifoliates. Unacceptable crop injury may occur to alfalfa seedlings less than the 2 trifoliolate leaf stage.

V-10137 1 EC Herbicide plus Buctril applications made when temperatures are expected to exceed 80°F at and 3 days following application can result in unacceptable crop injury. In the states not listed above, apply in the fall or spring to seedling alfalfa when the majority of the field has a minimum of 4 trifoliolate leaves. When alfalfa stand is uneven and conditions favor leafburn, unacceptable crop injury may occur to alfalfa in the 2 trifoliolate or smaller stage of growth. V-10137 1 EC Herbicide plus Buctril applications made when temperatures are expected to exceed 70°F at and 3 days following application can result in unacceptable crop injury. Crop leaf burn can occur following V-10137 1 EC Herbicide plus Buctril application. Warm, humid conditions may enhance leaf burn. New crop growth will not be affected.

⁽⁶⁾ **Restriction:** Do not apply when alfalfa is under moisture, temperature, insect or disease stress or has been stressed by other pesticide carryover or application.

USE DIRECTIONS FOR BUSHBERRY, CANEBERRY, POME FRUIT AND STONE FRUIT

Apply V-10137 1 EC Herbicide postemergence to actively growing grasses according to prescribed rates in Table 1, Crop Specific Use Directions, Restrictions and Limitations for V-10137 1 EC Herbicide. Crop injury to bushberry, caneberry, pome fruit and stone fruit can occur if V-10137 1 EC Herbicide is improperly applied. V-10137 1 EC Herbicide should not be applied directly over the top of these plant types. Instead spray should be directed at the base of the plant where grassy weeds are growing near the ground.

Restriction: Do not apply V-10137 1 EC Herbicide to bushberry, caneberry, pome fruit or stone fruit grown for root stock.

Non-bearing fruit and nut crops are plants which will not bear fruit or nuts for at least one year following V-10137 1 EC Herbicide application.

CANOLA (EXCEPT FLAX)

[Rapeseed Subgroup 20A (except flax seed, mustard seed and sesame seed)]

Table 3. REDUCED RATE V-10137 1 EC HERBICIDE TANK MIXES WITH BROADLEAF HERBICIDES (Refer to the use directions tables above for specific grasses and growth stages.)

PRODUCT	APPLICATION RATES/ACRE	ADJUVANT RECOMMENDATIONS	AMMONIUM SULFATE	
	ANNUAL GRASSES ⁽¹⁾		GROUND	AIR
V-10137 1 EC ⁽²⁾ Herbicide + Liberty ^{®(3)}	8 to 10 fl oz + Refer to Liberty label for use rate.	NIS at 0.25% v/v	3 lb/A	3 lb/A
V-10137 1 EC ⁽²⁾ Herbicide + Stinger ^{®(4)}	8 to 10 fl oz + Refer to Stinger label for use rate.	NIS at 0.25% v/v	3 lb/A	3 lb/A

⁽¹⁾ Annual grasses and sizes controlled with these tank mixtures are those that are identified in the DIRECTIONS FOR REDUCED RATE USE TO CONTROL SMALL ANNUAL GRASSES table.

⁽²⁾ **Restriction:** Do not apply V-10137 1 EC Herbicide tank mix during or after bolting or flowering or crop injury will occur.

⁽³⁾ For use only on Liberty Link Canola.

⁽⁴⁾ See Stinger label for weeds controlled.

COTTON (Including cotton grown for seed)

Table 4. V-10137 1 EC HERBICIDE TANK MIXED WITH COBRA HERBICIDE AND MSMA APPLIED POST DIRECTED TO COTTON

PRODUCT ⁽¹⁾	APPLICATION RATES/ACRE ⁽²⁾		CROP OIL CONCENTRATE ⁽³⁾ V/V	SPECIFIC USE INSTRUCTIONS
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	
V-10137 1 EC ⁽⁴⁾ Herbicide + Cobra + MSMA (4 lb/gal) or MSMA (6.6 lb/gal)	12 to 16 fl oz	16 to 32 fl oz	1%	Reduce broadcast rate in proportion to the band area actually treated.
	See Cobra Herbicide label for rates to control broadleaf weeds and height limitations for cotton. Refer to the V-10137 1 EC Herbicide label for weed height and species controlled.			
	See MSMA label for rates to control broadleaf weeds and height limitations for cotton. Refer to the V-10137 1 EC Herbicide label for weed height and species controlled.			

- ⁽¹⁾ Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.
- ⁽²⁾ If grass regrowth occurs or an additional flush of new grass emerges, make a second application of V-10137 1 EC Herbicide alone (without a tank mix herbicide), according to the appropriate size and rate prescribed.
- ⁽³⁾ Always use a crop oil concentrate at the listed rate (but not less than 1 pt/A) in the finished spray volume.
- ⁽⁴⁾ If at the time of application, grass height is so tall that post-directed applications cannot get good coverage over the top of the grassy weeds, then poor control may result and a second (non-post directed) application of V-10137 1 EC Herbicide may be necessary.

Table 5. V-101371 EC HERBICIDE TANK MIXED WITH GLYPHOSATE TO CONTROL EMERGED GRASSES IN COTTON AS A BROADCAST APPLICATION

PRODUCT	APPLICATION RATE/ACRE ⁽¹⁾		ADJUVANT		SPECIFIC USE INSTRUCTIONS
	ANNUAL GRASSES	PERENNIAL GRASSES	Glyphosate formulation with built in adjuvant	Glyphosate formulation without built in adjuvant	
V-101371 EC Herbicide + Glyphosate	9 to 16 fl oz	12 to 32 fl oz	Ammonium sulfate at 8.5 to 17 lb per 100 gals of carrier plus glyphosate label adjuvant recommendation.	Ammonium sulfate at 8.5 to 17 lb per 100 gals of carrier plus Non-ionic Surfactant (NIS) at 0.25% v/v.	See charts for grasses controlled. Use a minimum of 10 gals of spray solution per acre.
	See glyphosate label for rates to control broadleaf weeds and height limitations for cotton.				

⁽¹⁾ If grass regrowth occurs or an additional flush of new grass emerges, make a second application of V-101371 EC Herbicide at the prescribed rate with the appropriate amount of crop oil.

DRY AND SUCCULENT SHELLED BEANS
[Succulent Shelled Pea and Bean Subgroup 6B]
[Dried Shelled Pea and Bean (except Soybean 6C)]

Table 6. V-101371 EC HERBICIDE TANK MIXES WITH BROADLEAF HERBICIDES FOR DRY AND SUCCULENT SHELLED BEANS

(Refer to the use directions tables above for specific grasses and growth stages.)

PRODUCT ⁽¹⁾	APPLICATION RATES/ACRE ⁽²⁾		ADJUVANT	
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	AIR
V-101371 EC Herbicide + Basagran®	9 to 12 fl oz + Refer to Basagran label for use rate.	12 to 24 fl oz + Refer to Basagran label for use rate.	COC at 1% v/v + AMS at 2.5 lb/A	COC at 1% v/v + AMS at 17 lb/100 gal v/v
V-101371 EC Herbicide + Raptor	9 to 12 fl oz + Refer to Raptor label for use rate.	-	NIS at 0.25% v/v + AMS at 2.5 lb/A	NIS at 0.25% v/v + AMS at 17 lb/100 gal

⁽¹⁾ Broadleaf weed control may be reduced when grass populations are tall enough or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.

⁽²⁾ If grass regrowth occurs or an additional flush of new grass emerges, make a second application of V-101371 EC Herbicide alone (without a tank mix herbicide), according to the appropriate size and rate prescribed.

FLAX

Table 7. REDUCED RATE V-10137 1 EC HERBICIDE TANK MIXES WITH BROADLEAF HERBICIDES FOR FLAX

(Refer to the use directions tables above for specific grasses and growth stages.)

PRODUCT	APPLICATION RATES/ACRE	ADJUVANT	
	ANNUAL GRASSES ⁽¹⁾	GROUND	AIR
V-10137 1 EC Herbicide + Bronate Advanced™ ^(2, 3)	6 to 9 fl oz + Refer to Bronate Advanced label for use rate.	AMS at 2.4 to 4.0 lb/A + NIS at 0.125% v/v	AMS at 2.5 to 4.0 lb/A
V-10137 1 EC Herbicide + Bronate ^(2, 3)	6 to 9 fl oz + Refer to Bronate label for use rate.	AMS at 2.4 to 4.0 lb/A + NIS at 0.125% v/v	AMS at 2.5 to 4.0 lb/A
V-10137 1 EC Herbicide + Buctril ^(2, 3)	6 to 9 fl oz + Refer to Buctril label for use rate.	AMS at 2.4 to 4.0 lb/A + NIS at 0.125% v/v	AMS at 2.5 to 4.0 lb/A
V-10137 1 EC Herbicide + MCPA ^(2, 3)	8 to 10 fl oz + Refer to MCPA label for use rate.	AMS at 2.4 to 4.0 lb/A + NIS at 0.125% v/v	AMS at 2.5 to 4.0 lb/A
V-10137 1 EC Herbicide + Curtail® M ^(2, 3)	6 to 9 fl oz + Refer to Curtail M label for use rate.	AMS at 2.4 to 4.0 lb/A + NIS at 0.125% v/v	AMS at 2.5 to 4.0 lb/A

⁽¹⁾ Annual grasses and sizes controlled with these tank mixtures are those that are identified in the DIRECTIONS FOR REDUCED RATE USE TO CONTROL SMALL ANNUAL GRASSES table.

⁽²⁾ **Restriction:** Do not apply V-10137 1 EC Herbicide tank mix during or after the bud stage or to ornamental flax or crop injury may occur.

⁽³⁾ **Restriction:** Do not apply tank mixes if temperatures are expected to exceed 85°F at (or 3 days following) application or crop injury may occur.

PEANUT

Table 8. V-101371 EC HERBICIDE TANK MIXES WITH BROADLEAF HERBICIDES FOR PEANUT (Refer to the use directions tables above for specific grasses and growth stages.)

PRODUCT ⁽¹⁾	APPLICATION RATES/ACRE ⁽²⁾	ADJUVANT RECOMMENDATIONS	
	ANNUAL GRASSES	GROUND	AIR
V-101371 EC Herbicide + Basagran	9 to 16 fl oz + Refer to Basagran label for use rate.	COC at 1% v/v + AMS at 2.5 lb/A	COC at 1% v/v + AMS at 17 lb/100 gals
V-101371 EC Herbicide + Ultra Blazer®	9 to 16 fl oz + Refer to Ultra Blazer label for use rate.	COC at 1% v/v + AMS at 2.5 lb/A	COC at 1% v/v + AMS at 17 lb/100 gals
V-101371 EC Herbicide + Storm®	9 to 16 fl oz + Refer to Storm label for use rate.	COC at 1% v/v + AMS at 2.5 lb/A	COC at 1% v/v + AMS at 17 lb/100 gals

⁽¹⁾ Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.

⁽²⁾ If grass regrowth occurs or an additional flush of new grass emerges, make a second application of V-101371 EC Herbicide alone (without a tank mix herbicide), according to the appropriate size and rate prescribed.

USE DIRECTIONS FOR GRASS SUPPRESSION FOR HARVEST EFFICIENCY IN PEANUT WITH V-101371 EC HERBICIDE			
GRASS SPECIES	WEED STAGE	APPLICATION RATES	
		MINIMUM RATE fl oz/A	MAXIMUM RATE fl oz/A
Annual and perennial grasses that exceed height claimed for control on height charts “RECOMMENDATIONS FOR ANNUAL GRASSES” and “RECOMMENDATIONS FOR PERENNIAL GRASSES”	Up to and including grasses in the seed head stage	32	64
<ul style="list-style-type: none"> • Restriction: Do not apply as part of a tank mix when applying V-101371 EC Herbicide for grass suppression. • Add a crop oil concentrate at 1 qt/A by ground to the finished spray volume. 			

SOYBEAN

Table 9. V-10137 1 EC Herbicide TANK MIX WITH BROADLEAF HERBICIDES FOR THE CONTROL OF VOLUNTEER CORN (INCLUDING ROUNDUP READY) IN SOYBEAN
(Refer to the use directions tables above for specific volunteer corn sizes and use rates.)

PRODUCT	WEED SIZE AND APPLICATION RATES		SPRAY ADDITIVES			
	VOLUNTEER CORN HEIGHT (inches)	V-10137 1 EC HERBICIDE RATES/ACRE	GROUND APPLICATION		AIR APPLICATION	
			NIS	AMS	NIS	AMS
V-10137 1 EC Herbicide + glyphosate ^(1,2,3) Refer to glyphosate label for use rate. (Roundup Ready soybeans only)	Up to 12 Up to 24 Up to 36	6 fl oz 9 fl oz 12 fl oz	Adjuvant Loaded Glyphosate: None Required Adjuvant Unloaded Glyphosate: NIS at 0.25% v/v	8.5 to 17 lb/100 gals of spray solution	Adjuvant Loaded Glyphosate: None Required Adjuvant Unloaded Glyphosate: NIS at 0.25% v/v	8.5 to 17 lb/100 gals of spray solution
V-10137 1 EC Herbicide + Firstrate® Refer to Firstrate label for use rate.	Up to 12 Up to 24 Up to 36	6 fl oz 9 fl oz 12 fl oz	NIS at 0.25% v/v	AMS at 2.5 lb/A	NIS at 0.25% v/v	17 lb/100 gals of spray solution
V-10137 1 EC Herbicide + Pursuit 70 DG Refer to Pursuit 70 DG label for use rate.	Up to 12 Up to 24 Up to 36	6 fl oz 9 fl oz 12 fl oz	NIS at 0.25% v/v	AMS at 2.5 lb/A	NIS at 0.25% v/v	17 lb/100 gals of spray solution
V-10137 1 EC Herbicide + Raptor Refer to Raptor label for use rate.	Up to 12 Up to 24 Up to 36	6 fl oz 9 fl oz 12 fl oz	NIS at 0.25% v/v	AMS at 2.5 lb/A	NIS at 0.25% v/v	17 lb/100 gals of spray solutions

- ⁽¹⁾ This tank mix may be applied postemergence to Roundup Ready soybeans up through the full flowering stage. **Restriction:** Do not apply less than 60 days before harvest.
- ⁽²⁾ Avoid contact with foliage, green stems or fruit crops or any desirable plants and trees, other than soybeans with the Roundup Ready gene as severe plant injury or death will result.
- ⁽³⁾ **Restriction:** Do not allow the V-10137 1 EC Herbicide plus glyphosate to mist, drip, drift or splash onto desirable vegetation as minute quantities of the tank mix can cause severe damage or destruction to the crops, plants or other areas on which treatment was not intended. The likelihood of injury occurring from drift of this product is greatest when winds are gusty or in excess of 5 miles per hour. Even under lesser wind velocities, avoid conditions that allow spray drift to occur such as combinations of spray pressure and nozzle type that will result in fine particles (mist) that are likely to drift.

Table 10. V-101371 EC HERBICIDE TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEAN
(Refer to the use directions tables above for specific grasses and growth stages.)

PRODUCT ⁽¹⁾	APPLICATION RATES/ACRE ⁽²⁾	SPRAY ADDITIVE RECOMMENDATIONS			
	ANNUAL GRASSES ⁽³⁾	GROUND APPLICATION		AIR APPLICATION	
		COC/NIS ⁽⁴⁾	AMS	COC/NIS ⁽⁴⁾	AMS
V-101371 EC Herbicide + Cobra (EPA Reg. No. 59639-34)	9 to 20 fl oz + 6 to 12 fl oz	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1 to 2 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v plus COC at 0.25% v/v or COC 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-101371 EC Herbicide + Firstrate ⁽⁵⁾	9 to 20 fl. oz + Refer to Firstrate label for use rate.	NIS at 0.25% v/v or COC at 1 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-101371 EC Herbicide + Flexstar® HL ⁽⁵⁾	9 to 20 fl oz + Refer to the Flexstar HL label for use rate.	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1 to 2 pt/A	AMS at 2.5 lb/A	NIS at 0.25% plus COC at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-101371 EC Herbicide + Frontrow™ ⁽⁵⁾	9 to 20 fl oz + Refer to Frontrow label for use rate.	NIS at 0.25% v/v or COC at 1 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-101371 EC Herbicide + Harmony® GT XP ⁽⁵⁾	9 to 12 fl oz + Refer to Harmony GT XP label for use rate.	Nis at 0.125 to 0.25% v/v	AMS at 2.5 lb/A	-	-
V-101371 EC Herbicide + Phoenix™ (EPA Reg. No. 59639-118)	9 to 20 fl oz + 6 to 12.5 fl oz	NIS at 0.25% v/v plus COC at 0.125 to 0.25% v/v or COC at 1 pt/A	AMS at 2.5 lb/A	NIS at 0.25% plus COC at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-101371 EC Herbicide + Pursuit 70 DG ⁽⁵⁾	12 to 20 fl oz + Refer to Pursuit 70 DG label for use rate.	NIS at 0.25% v/v or COC at 1 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-101371 EC Herbicide + Raptor (1 AS) ⁽⁵⁾	12 to 20 fl oz + Refer to Raptor label for use rate.	NIS at 0.25% v/v or COC at 1 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-101371 EC Herbicide + Resource® (EPA Reg. No. 59639-82)	9 to 20 fl oz + 4 to 12 fl oz	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1 to 2 pt/A	AMS at 2.5 lb/A	-	-

**Table 10. V-10137 1 EC HERBICIDE TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEAN
(Refer to the use directions tables above for specific grasses and growth stages.)**

PRODUCT ⁽¹⁾	APPLICATION RATES/ACRE ⁽²⁾	SPRAY ADDITIVE RECOMMENDATIONS			
	ANNUAL GRASSES ⁽³⁾	GROUND APPLICATION		AIR APPLICATION	
		COC/NIS ⁽⁴⁾	AMS	COC/NIS ⁽⁴⁾	AMS
V-10137 1 EC Herbicide + Cobra (EPA Reg. No. 59639-34) + Firstrate ⁽⁵⁾	9 to 20 fl oz + 6 to 12.5 fl oz + Refer to Firstrate label use rate.	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1 to 2 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-10137 1 EC Herbicide + Cobra (EPA Reg. No. 59639-34) + Harmony GT XP ⁽⁵⁾	9 to 12 fl oz + 6 to 8 fl oz + Refer to Harmony GT XP label for use rate.	NIS at 0.125 to 0.25% v/v plus COC at 0.125% v/v	AMS at 2.5 lb/A	-	-
V-10137 1 EC Herbicide + Cobra (EPA Reg. No. 59639-34) + Pursuit 70 DG ⁽⁵⁾	12 to 20 fl oz + 6 to 12.5 fl oz + Refer to Pursuit 70 DG label for use rate.	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1 to 2 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-10137 1 EC Herbicide + Cobra (EPA Reg. No. 59639-34) + Raptor (1 AS) ⁽⁵⁾	12 to 20 fl oz + 6 to 12.5 fl oz + Refer to Raptor (1 AS) label for use rate.	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1 to 2 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-10137 1 EC Herbicide + Cobra (EPA Reg. No. 59639-34) + Resource (EPA Reg. No. 59639-82)	9 to 20 fl oz + 6 to 12.5 fl oz + 4 to 6 fl oz	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1 to 2 pt/A	AMS at 2.5 lb/A	-	-
V-10137 1 EC Herbicide + Firstrate + Flexstar HL ⁽⁵⁾	9 to 20 fl oz + Refer to Firstrate and Flexstar HL labels for use rates.	NIS at 0.25% v/v plus COC at 0.25% v/v or Equivalent blended product or COC at 1 to 2 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution

**Table 10. V-10137 1 EC HERBICIDE TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEAN
(Refer to the use directions tables above for specific grasses and growth stages.)**

PRODUCT ⁽¹⁾	APPLICATION RATES/ACRE ⁽²⁾	SPRAY ADDITIVE RECOMMENDATIONS			
	ANNUAL GRASSES ⁽³⁾	GROUND APPLICATION		AIR APPLICATION	
		COC/NIS ⁽⁴⁾	AMS	COC/NIS ⁽⁴⁾	AMS
V-10137 1 EC Herbicide + <i>Phoenix</i> (EPA Reg. No. 59639-118) + Firstrate ⁽⁵⁾	12 to 20 fl oz + 6 to 12.5 fl oz + Refer to Firstrate label for use rate.	NIS at 0.25% v/v plus COC at 0.125 to 0.25% v/v or COC at 1 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-10137 1 EC Herbicide + <i>Phoenix</i> (EPA Reg. No. 59639-118) + Pursuit 70 DG ⁽⁵⁾	16 to 20 fl oz + 6 to 12.5 fl oz + Refer to Pursuit 70 DG label for use rate.	NIS at 0.25% v/v plus COC at 0.125 to 0.25% v/v or COC at 1 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-10137 1 EC Herbicide + <i>Phoenix</i> (EPA Reg. No. 59639-118) + Raptor (1 AS) ⁽⁵⁾	12 to 20 fl oz + 6 to 12.5 fl oz + Refer to Raptor (1 AS) label for use rate.	NIS at 0.25% v/v plus COC at 0.125 to 0.25% v/v or COC at 1 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-10137 1 EC Herbicide + <i>Phoenix</i> (EPA Reg. No. 59639-118) + <i>Resource</i> (EPA Reg. No. 59639-82)	9 to 20 fl oz + 6 to 12.5 fl oz + 4 to 6 fl oz	NIS at 0.25% v/v plus COC at 0.125 to 0.25% v/v or COC at 1 pt/A	AMS at 2.5 lb/A	-	-
V-10137 1 EC Herbicide + <i>Resource</i> (EPA Reg. No. 59639-82) + Pursuit 70 DG ⁽⁵⁾	12 to 20 fl oz + 4 fl oz + Refer to Pursuit 70 DG label for use rate.	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1 to 2 pt/A	AMS at 2.5 lb/A	-	-
V-10137 1 EC Herbicide + Synchrony XP (mp) ⁽⁵⁾	12 to 20 fl oz + Refer to Synchrony XP (mp) label for use rate.	NIS at 0.25% v/v or COC at 1 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution

**Table 10. V-10137 1 EC HERBICIDE TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEAN
(Refer to the use directions tables above for specific grasses and growth stages.)**

PRODUCT ⁽¹⁾	APPLICATION RATES/ACRE ⁽²⁾	SPRAY ADDITIVE RECOMMENDATIONS			
	ANNUAL GRASSES ⁽³⁾	GROUND APPLICATION		AIR APPLICATION	
		COC/NIS ⁽⁴⁾	AMS	COC/NIS ⁽⁴⁾	AMS
V-10137 1 EC Herbicide + Synchrony XP (mp) ⁽⁵⁾ (STS Soybeans Only)	12 to 20 fl oz + Refer to Synchrony XP (mp) (STS Soybeans Only) label for use rate.	NIS at 0.25% v/v or COC at 1 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-10137 1 EC Herbicide + Cobra (EPA Reg. No. 59639-34) + Resource (EPA Reg. No. 59639-82) + Firstrate ⁽⁵⁾	9 to 12 fl oz + 6 to 12.5 fl oz + 4 to 6 fl oz + Refer to Firstrate label for use rate.	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1 to 2 pt/A	AMS at 2.5 lb/A	-	-
V-10137 1 EC Herbicide + Cobra (EPA Reg. No. 59639-34) + Synchrony XP (mp) ⁽⁵⁾	12 to 20 fl oz + 6 to 12 fl oz + Refer to Synchrony XP (mp) label for use rate.	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1 to 2 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-10137 1 EC Herbicide + Cobra (EPA Reg. No. 59639-34) + Synchrony XP (mp) ⁽⁵⁾ (STS Soybeans Only)	12 to 20 fl oz + 6 to 12 fl oz + Refer to Synchrony XP (mp) (STS Soybeans Only) label for use rate.	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1 to 2 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution

**Table 10. V-10137 1 EC HERBICIDE TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEAN
(Refer to the use directions tables above for specific grasses and growth stages.)**

PRODUCT ⁽¹⁾	APPLICATION RATES/ACRE ⁽²⁾	SPRAY ADDITIVE RECOMMENDATIONS			
	ANNUAL GRASSES ⁽³⁾	GROUND APPLICATION		AIR APPLICATION	
		COC/NIS ⁽⁴⁾	AMS	COC/NIS ⁽⁴⁾	AMS
V-10137 1 EC Herbicide + <i>Phoenix</i> (EPA Reg. No. 59639-118) + <i>Resource</i> (EPA Reg. No. 59639-82) + Firstrate ⁽⁵⁾	9 to 20 fl oz + 6 to 12.5 fl oz + 4 to 6 fl oz + Refer to Firstrate label for use rates.	NIS at 0.25% v/v plus COC at 0.125 to 0.25% v/v or COC at 1 pt/A	AMS at 2.5 lb/A	-	-
V-10137 1 EC Herbicide + <i>Phoenix</i> (EPA Reg. No. 59639-118) + Synchrony XP (mp) ⁽⁵⁾	12 to 20 fl oz + 6 to 12 fl oz + Refer to Synchrony XP (mp) label for use rate.	NIS at 0.25% v/v plus COC at 0.125 to 0.25% v/v or COC at 1 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution
V-10137 1 EC Herbicide + <i>Phoenix</i> (EPA Reg. No. 59639-82) + Synchrony XP (mp) ⁽⁵⁾ (STS Soybeans Only)	12 to 20 fl oz + 6 to 12 fl oz + Refer to Synchrony XP (mp) (STS Soybeans Only) label for use rate.	NIS at 0.25% v/v plus COC at 0.125 to 0.25% v/v or COC at 1 pt/A	AMS at 2.5 lb/A	NIS at 0.25% v/v plus COC at 0.25% v/v or COC at 1% v/v (but not less than 1 pt/A)	17 lb/100 gals of spray solution

- ⁽¹⁾ Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.
- ⁽²⁾ If grass regrowth occurs or an additional flush of new grass emerges, make a second application of V-10137 1 EC Herbicide alone (without a tank mix herbicide), according to the appropriate size and rate prescribed.
- ⁽³⁾ Annual grasses and sizes controlled with these tank mixtures are those that are identified in the **RECOMMENDATIONS FOR ANNUAL GRASSES** table.
- ⁽⁴⁾ Contact local Valent U.S.A. representative for proper COC/NIS adjuvant selection.
- ⁽⁵⁾ Refer to Firstrate, Flexstar HL, Frontrow, Harmony GT XP, Pursuit DG, Raptor and Synchrony XP (mp) for geographic and rotational restrictions.

SUGAR BEET

Table 11. V-10137 1 EC HERBICIDE TANK MIXED WITH BROADLEAF SUGAR BEET HERBICIDES

PRODUCTS	APPLICATION RATES/A	ADJUVANT INFORMATION
V-10137 1 EC Herbicide	9 to 12 fl oz	
+		
Betamix®	Refer to label for use rate.	None required
or		
Betanex®	Refer to label for use rate.	None required
or		
Progress®	Refer to label for use rate.	None required
and/or		
Stinger	Refer to label for use rate.	See below
and/or		
Upbeet®	Refer to label for use rate.	See below
NIS at 0.25% unless Betamix, Betanex or Progress is in the tank, then use no adjuvant.		

TABLE 12. V-10137 1 EC HERBICIDE PLUS BETANEX OR BETAMIX TANK MIX FOR THREE SEQUENTIAL APPLICATIONS FOR ANNUAL GRASS CONTROL (MICRO RATE APPLICATION)

PRODUCT	APPLICATION RATES/ACRE ⁽¹⁾	GRASSES CONTROLLED (inches)	METHYLATED SEED OIL ⁽²⁾	
	ANNUAL GRASSES		GROUND	AIR
V-10137 1 EC Herbicide	3 to 6 fl oz	Green Foxtail (1 to 2) Yellow Foxtail (1 to 2) Barnyardgrass (1 to 2) Wild Oat (1 to 2) Volunteer Cereals (1 to 2)	1.5% v/v	1.5% v/v
+				
Betanex	Refer to label for use rate.			
or				
Betamix	Refer to label for use rate.			
or				
Progress	Refer to label for use rate.			
or				
Stinger	Refer to label for use rate.			
or				
UpBeet	Refer to label for use rate.			

⁽¹⁾ Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.

⁽²⁾ Always use a methylated seed oil at the listed rate (but not less than 1 pt/A) in the finished spray volume.

Directions for Use for Micro-Rate Applications to Sugar Beet

Multiple micro-rate applications of V-10137 1 EC Herbicide in tank mixtures with reduced rates of Betanex or Betamix and methylated seed oils may be applied by air or ground equipment to sugar beet to control early germinating annual grasses listed above. All use precautions and restrictions on the Betanex and Betamix master labels must be followed.

Table 13. TANK MIX APPLICATION OF V-101371 EC HERBICIDE AND FUNGICIDES FOR CONTROL OF GRASS WEEDS AND DISEASES IN SUGAR BEET

PRODUCT ⁽¹⁾	APPLICATION RATES/ACRE ⁽²⁾		ADJUVANT
	ANNUAL GRASSES	PERENNIAL GRASSES	
V-101371 EC Herbicide + Eminent [®]	9 to 12 fl oz + Refer to Eminent label for use rate.	12 to 24 fl oz + Refer to Eminent label for use rate.	NIS at 0.25% v/v
V-101371 EC Herbicide + Headline [®]	9 to 12 fl oz + Refer to Headline label for use rate.	12 to 24 fl oz + Refer to Headline label for use rate.	NIS at 0.25% v/v
V-101371 EC Herbicide + Gem [™]	9 to 12 fl oz + Refer to Gem label for use rate.	12 to 24 fl oz + Refer to Gem label for use rate.	NIS at 0.25% v/v

⁽¹⁾ Refer to V-101371 EC Herbicide and fungicide label for rates and weeds and diseases controlled.

⁽²⁾ If grass regrowth occurs, or an additional flush of new grass emerges, make a second application of V-101371 EC Herbicide alone (without a tank mix fungicide) according to the appropriate size and rate prescribed.

Table 14. TANK MIX APPLICATION OF V-101371 EC HERBICIDE AND INSECTICIDES FOR CONTROL OF GRASS WEEDS AND INSECTS IN ALFALFA, COTTON, MINT, PEANUT (INCLUDING PERENNIAL), SOYBEAN AND SUNFLOWER

PRODUCT ⁽¹⁾	APPLICATION RATES/ACRE ⁽²⁾		ADJUVANT RECOMMENDATION	CROPS					
	ANNUAL GRASSES	PERENNIAL GRASSES		Alfalfa ⁽³⁾	Cotton	Mint ^(3,4)	Peanut	Soybean	Sunflower
V-101371 EC Herbicide + Asana [®] XL	9 to 12 fl oz + Refer to Asana XL label for use rate.	12 to 24 fl oz + Refer to Asana XL label for use rate.	NIS at 0.25% v/v + AMS at 2.5 lb/A					X	X
V-101371 EC Herbicide + Baythroid [®]	9 to 12 fl oz + Refer to Baythroid label for use rate.	12 to 24 fl oz + Refer to Baythroid label for use rate.	NIS at 0.25% v/v + AMS at 2.5 lb/A	X				X	
V-101371 EC Herbicide + Danitol [®] 2.4 EC (EPA Reg. No. 59639-35)	9 to 12 fl oz + 10 2/3 to 16 fl oz	12 to 24 fl oz + 10 2/3 to 16 fl oz	NIS at 0.25% v/v + AMS at 2.5 lb/A		X		X		
V-101371 EC Herbicide + dimethoate	9 to 12 fl oz + Refer to dimethoate label for use rate.	12 to 24 fl oz + Refer to dimethoate label for use rate.	NIS at 0.25% v/v + AMS at 2.5 lb/A	X				X	

Table 14. TANK MIX APPLICATION OF V-101371 EC HERBICIDE AND INSECTICIDES FOR CONTROL OF GRASS WEEDS AND INSECTS IN ALFALFA, COTTON, MINT, PEANUT (INCLUDING PERENNIAL), SOYBEAN AND SUNFLOWER

V-101371 EC Herbicide + Lorsban®	9 to 12 fl oz + Refer to Lorsban label for use rate.	12 to 24 fl oz + Refer to Lorsban label for use rate.	NIS at 0.25% v/v + AMS at 2.5 lb/A	X				X	
V-101371 EC Herbicide + Orthene® 75 S or Orthene 97	9 to 12 fl oz + Refer to Orthene 75 S or Orthene 97 labels for use rates.	12 to 24 fl. oz + Refer to Orthene 75 S or Orthene 97 labels for use rates.	NIS at 0.25% v/v + AMS at 2.5 lb/A		X	X	X	X	
V-101371 EC Herbicide + Orthene 90 S	9 to 12 fl oz + Refer to Orthene 90 S label for use rate.	12 to 24 fl oz + Refer to Orthene 90 S label for use rate.	NIS at 0.25% v/v + AMS at 2.5 lb/A		X	X	X	X	
V-101371 EC Herbicide + Pounce®	9 to 12 fl oz + Refer to Pounce label for use rate.	12 to 24 fl oz + Refer to Pounce label for use rate.	NIS at 0.25% v/v + AMS at 2.5 lb/A	X				X	
V-101371 EC Herbicide + Warrior®	9 to 12 fl oz + Refer to Warrior label for use rate.	12 to 24 fl oz + Refer to Warrior label for use rate.	NIS at 0.25% v/v + AMS at 2.5 lb/A					X	X

⁽¹⁾ Refer to V-101371 EC Herbicide and insecticide label for rates and weeds and insects controlled.

⁽²⁾ If grass regrowth occurs, or an additional flush of new grass emerges, make a second application of V-101371 EC Herbicide alone (without a tank mix insecticide) according to the appropriate size and rate prescribed.

⁽³⁾ Certain insecticides may cause temporary phytotoxic symptoms on alfalfa and mint foliage. Refer to the insecticide label for further information. It is suggested that prior to using any of these insecticide/herbicide tank mixtures, that a small area of the field be treated first and observations for crop injury be made prior to treating the whole field.

⁽⁴⁾ The V-101371 EC Herbicide rate should be 9 to 12 fl oz/A for annual grass control in baby mint, minimum of 12 fl oz/A for annual grass control in established mint and 16 to 32 fl oz/A for perennial grass control.

FALLOW LAND

DIRECTIONS FOR USE

V-10137 1 EC Herbicide may be used to control annual and perennial grasses in land that has been left fallow the previous year and other non-producing agricultural areas. Apply V-10137 1 EC Herbicide at 12 to 16 fl oz/A for annual grasses and 16 to 32 fl oz/A for perennial grasses. When both grass and broadleaf weeds are the target pest, V-10137 1 EC Herbicide may be tank mixed with 2,4-D ester or Banvel® SGF for broad spectrum control. When both annual and perennial grasses occur in the same field, use a minimum of 16 fl oz/A V-10137 1 EC Herbicide rate.

Precautions

- Use a minimum spray volume of 5 gals/A for aerial applications and 15 gals/A for ground applications. Apply only to actively growing grasses when the first grass reaches the recommended weed height as specified by the Recommendations for Annual and Perennial Grasses section of this label.
- Annual grasses that emerge after the V-10137 1 EC Herbicide application will not be controlled, and a second application may be necessary.
- The control of perennial grasses may require more than 1 application in non-tilled areas.

Restrictions

- Do not apply more than 32 fl oz/A (0.242 lb ai/A) V-10137 1 EC Herbicide per application.
- Do not apply more than 64 fl oz/A (0.485 lb ai/A) V-10137 1 EC Herbicide per year.
- Do not apply to grasses that have tillered, formed seedheads or exceeded recommended growth stage.
- Do not use flood jet nozzles.
- Do not apply to drought stressed grasses.
- Do not mow area for 2 weeks prior to or after the V-10137 1 EC Herbicide application.

TABLE 15. V-10137 1 EC HERBICIDE IN TANK MIXES TO CONTROL ANNUAL AND PERENNIAL GRASSES IN FALLOW LAND

PRODUCT	APPLICATION RATES/ACRE ⁽¹⁾		ADJUVANT RECOMMENDATION	
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	AIR
V-10137 1 EC Herbicide + 2,4-D ester or Banvel SGF	12 to 16 fl oz + See 2,4-D ester or Banvel SGF labels for use rates.	16 to 32 fl oz + See 2,4-D ester or Banvel SGF labels for use rates	NIS at 0.25% v/v or COC at 1% + AMS at 2.5 lb/A	NIS at 0.25% v/v or COC at 1% + AMS at 17 lb/100 gals

⁽¹⁾ Refer to V-10137 1 EC Herbicide label for weed height and species control. Review Banvel SGF and 2,4-D labels for crop restrictions, use rates and weeds controlled.

USE DIRECTIONS FOR GRASS SUPPRESSION IN NON-CROP AREAS WITH V-10137 1 EC HERBICIDE

GRASS SPECIES	WEED STAGE	APPLICATION RATES	
		MINIMUM RATE fl oz/A	MAXIMUM RATE fl oz/A
Annual and perennial grasses that exceed height claimed for control on height chart above.	Up to and including grasses in the seed head stage	24	32
<ul style="list-style-type: none"> • Restriction: Do not apply as part of a tank mix when applying V-10137 1 EC Herbicide for grass suppression. • Add a crop oil concentrate at 1 qt/A by ground to the finished spray volume. 			

TABLE 16. V-10137 1 EC HERBICIDE FOR THE CONTROL AND/OR SUPPRESSION OF TALL FESCUE IN NATIVE PRAIRIE WARM-SEASON GRASS RESTORATION PROJECTS

PRODUCT	PRODUCT RATE	GRASS WEEDS CONTROLLED/SUPPRESSED		WEED STAGE
		Common Name	Scientific Name	
V-10137 1 EC Herbicide	12 to 16 fl oz/A	Tall Fescue	<i>Festuca arundinacea</i>	4 to 6 inches tall (40 to 60% green-up)
<p>Adjuvant: V-10137 1 EC Herbicide must be applied with non-ionic surfactant at 0.25% v/v, plus a spray grade ammonium sulfate at 2.5 to 4 lb/A.</p> <p>Recommended Mixing Order: Thoroughly mix spray grade ammonium sulfate in water, add V-10137 1 EC Herbicide, then add non-ionic surfactant.</p>				

SPECIAL APPLICATION PRECAUTIONS AND RESTRICTIONS
<p>Burn or mow fields a minimum of 3 weeks prior to application to remove excess crop residue. Apply in the spring, at 40 to 60% tall fescue green-up, prior to emergence of warm-season grasses. Do not mow area for 2 weeks after the V-10137 1 EC Herbicide application.</p> <p>Apply in a minimum of 15 to 20 gals of water per acre at a spray pressure of 40 to 60 PSI at the nozzle. Apply using flat fan or hollow cone nozzles. Do not use flood jet nozzles.</p> <p>Apply only to fields that have warm-season grasses established for 2 years. Applications of V-10137 1 EC Herbicide to emerged warm-season grasses may cause injury. Do not apply to warm-season grasses grown for seed.</p> <p>Restriction: Do not graze treated fields or feed treated forage and or hay to livestock.</p> <p>NOTE: V-10137 1 EC Herbicide applications are most effective if applied when average nighttime temperatures are consistently greater than or equal to 47 degrees Fahrenheit.</p>

TABLE 17. V-10137 1 EC HERBICIDE FOR THE SUPPRESSION OF TALL FESCUE SEED-HEADS IN NON-PRODUCING AGRICULTURAL AREAS

PRODUCT	PRODUCT RATES	SUPPRESSION	APPLICATION TIMING
V-10137 1 EC Herbicide	3 to 4 fl oz/A	Tall Fescue Seed-Heads (<i>Festuca arundinacea</i>)	(50 to 90% Tall Fescue green-up in the spring) or 3 weeks prior to dormancy in the fall.
<p>ADJUVANT: V-10137 1 EC Herbicide must be applied with crop oil concentrate at 1 qt/A, plus a spray grade ammonium sulfate at 2.5 to 4 lb/A.</p> <p>Recommended Mixing Order: Thoroughly mix spray grade ammonium sulfate in water, add V-10137 1 EC Herbicide, then add crop oil concentrate.</p> <p>Note: Use crop oil concentrate at 2 pt/A with fall applications.</p>			

SPECIAL APPLICATION PRECAUTIONS AND RESTRICTIONS

- Apply at 50 to 90% tall fescue green-up.
- Use the higher *V-10137 1* EC Herbicide rate if less tall fescue green matter is present.
- Apply in a minimum of 15 to 20 gals of water per acre at a spray pressure of 40 to 60 psi at the nozzle. Apply using flat fan or hollow cone nozzles.
- 2,4-D ester, Tordon[®] 22K, Grazon[®] P+D or Crossbow[®] maybe added to this tank mix for broadleaf control (see 2,4-D ester label for weeds controlled)

Restrictions

- Do not mow area for 2 weeks after the *V-10137 1* EC Herbicide application.
- Do not use flood nozzles.
- Do not graze treated fields or feed treated forage and or hay to livestock

DIRECTIONS FOR USE IN ORNAMENTALS

For ornamental plant uses, V-101371 EC Herbicide can be used to control labeled grass weeds in greenhouses, lathhouses, shadehouses, and around outdoor ornamentals, including nurseries, parks, roadside plantings, and structure landscapes.

IMPORTANT

V-101371 EC Herbicide successfully controls weeds in newly transplanted and established non-grassy ornamentals. Plant tolerance to V-101371 EC Herbicide at labeled rates has been found to be acceptable for the indicated genera and species listed below. Due to variability within species, crop growth stage, environmental conditions, and application techniques, it is recommended that the user determine if herbicide can be used safely on a few plants prior to widespread application. Neither the seller nor the manufacturer of V-101371 EC Herbicide have investigated the safety factor to ornamental plants not listed on the label.

The following plants have shown a tolerance for V-101371 EC Herbicide applications:

ORNAMENTAL TREES

COMMON NAME	SCIENTIFIC NAME
Alder, Red	<i>Alnus rubra</i>
Ash	<i>Fraxinus</i> spp.
Basswood	<i>Tilia</i> spp.
Birch, European White	<i>Betula pendula</i>
Birch, River	<i>Betula nigra</i>
Birch, White	<i>Betula papyrifera</i>
Crabapple, Flowering	<i>Malus halliana</i>
Dogwood, Flowering	<i>Cornus florida</i>
Goldon Chain Tree	<i>Laburnum anagyroides</i>
Maples	<i>Acer</i> spp.
Mulberry, White	<i>Morus alba</i>
Oaks	<i>Quercus</i> spp.
Olive, Wild	<i>Elaeagnus angustifolia</i>
Redbud, Eastern	<i>Cercis canadensis</i>
Sweet Gum, American	<i>Liquidambar styraciflua</i>

GROUND COVERS

COMMON NAME	SCIENTIFIC NAME
Bugleweed, Carpet	<i>Ajuga reptans</i>
Ivy, English	<i>Hedera helix</i>
Japanese Spurge	<i>Pachysandra terminalis</i>
Lilyturf	<i>Liriope muscari</i>
Moneywort	<i>Lysimachia nummularia</i>
Mondo Grass, White	<i>Ophiopogon jaburan</i>
Mondo Grass Dwarf	<i>Ophiopogon japonicus</i>
Periwinkle, Lesser	<i>Vinca minor</i>

GARDEN FLOWERS AND PLANTS

COMMON NAME	SCIENTIFIC NAME
Ageratum	<i>Ageratum</i> spp.
Alyssum*, Sweet	<i>Lobularia maritima</i>
Asparagus Fern	<i>Asparagus setaceus</i>
Bleeding Heart	<i>Dicentra spectabilis</i>
Cast Iron Plant	<i>Aspidistra elatior</i>
Chrysanthemum	<i>Chrysanthemum</i> spp.
Cinquefoil	<i>Potentilla</i> spp.
Coleus	<i>Coleus</i> spp.
Coralbells	<i>Heuchera sanguinea</i>
Cranesbill	<i>Geranium</i> spp.
Dahlia	<i>Dahlia</i> spp.
Daisy, Trailing African	<i>Osteospermum fruticosum</i>
Daylily	<i>Hemerocallis</i> spp.
Dusty Miller	<i>Senecio cineraria</i>
Euonymus	<i>Euonymus</i> spp.
Gazania	<i>Gazania</i> spp.
Geranium, House	<i>Pelargonium hortorum</i>
Heather, False	<i>Cuphea hyssopifolia</i>
Hosta	<i>Hosta fortunei</i>
Iris	<i>Iris</i> spp.
Jasmine Tobacco	<i>Nicotiana alata</i>
Loosestrife	<i>Lythrum salicaria</i>
Marigold	<i>Tagetes</i> spp.
Partridgeberry	<i>Mitchella repens</i>
Petunia*	<i>Petunia hybrida</i>
Phlox	<i>Phlox</i> spp.
Pinks	<i>Dianthus</i> spp.
Portulaca	<i>Portulaca grandiflora</i>
Salvia	<i>Salvia</i> spp.
Saxifrage	<i>Saxifraga</i> spp.
Sedum	<i>Sedum</i> spp.
Selloum	<i>Philodendron selloum</i>
Snapdragon*	<i>Antirrhinum majus</i>
Sweet Flag	<i>Acorus gramineus</i>
Tickseed	<i>Coreopsis grandiflora</i>
Touch-Me-Not	<i>Impatiens</i> spp.
Verbena	<i>Verbena</i> spp.
Violet	<i>Viola</i> spp.
Yarrow, Common	<i>Achillea millefolium</i>
Zinnia	<i>Zinnia elegans</i>

*Slight foliage or flower speckling has been observed on these species.

SHRUBS

COMMON NAME	SCIENTIFIC NAME
Abelia	<i>Abelia</i> spp.
Anise, Purple	<i>Illicium floridanum</i>
Aucuba	<i>Aucuba</i> spp.
Azalea*	<i>Rhododendron</i> spp.
Bamboo	<i>Bambusa</i> spp.
Barberry, Japanese	<i>Berberis thunbergii</i>
Barberry, Magellan	<i>Berberis buxifolia</i>
Bayberry	<i>Myrica pensylvanica</i>
Bottlebrush	<i>Callistemon citrinus</i>
Boxwood, Common	<i>Buxus sempervirens</i>
Camellia, Common	<i>Camellia japonica</i>
Candytuft	<i>Iberis sempervirens</i>
Cleyera	<i>Cleyera japonica</i>
Coralberry	<i>Ardisia crenata</i>
Crape Myrtle	<i>Lagerstroemia indica</i>
Coyote Brush	<i>Baccharis pilularis</i>
Fig, Creeping	<i>Ficus pumila</i>
Gardenia	<i>Gardenia</i> spp.
Holly	<i>Ilex</i> spp.
Honeysuckle	<i>Lonicera</i> spp.
Indian Hawthorn	<i>Raphiolepis indica</i>
Jasmine	<i>Jasminum</i> spp.
Jasmine, Asiatic	<i>Trachelospermum asiaticum</i>
Jasmine, Star	<i>Trachelospermum jasminoides</i>
Juniper	<i>Juniperus</i> spp.
Lantana	<i>Lantana</i> spp.
Nandina* Bamboo, Heavenly	<i>Nandinia domestica</i>
Oleander, Common	<i>Nerium oleander</i>
Oregon Grape	<i>Mahonia aquifolium</i>
Photinia	<i>Photinia</i> spp.
Pittosporum	<i>Pittosporum</i> spp.
Podocarpus	<i>Podocarpus</i> spp.
Privet	<i>Ligustrum</i> spp.
Pyracantha	<i>Pyracantha</i> spp.
Rhododendron	<i>Rhododendron</i> spp.
Rose	<i>Rosa</i> spp.
Spirea	<i>Spiraea bumalda</i>
Sweet Olive	<i>Osmanthus fragrans</i>
Viburnum	<i>Viburnum tinus</i>
Wisteria	<i>Wisteria</i> spp.
Yellow Sage/Shrub Verbena	<i>Lantana camara</i>

*Slight foliage or flower speckling has been observed on these species.

USE DIRECTIONS FOR ANNUAL GRASSES IN ORNAMENTALS

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

GRASS SPECIES	SCIENTIFIC NAME	WEED* HEIGHT INCHES	APPLICATION RATES	
			MINIMUM RATE fl oz/A ⁽¹⁾	MAXIMUM RATE ⁽²⁾
Barnyardgrass	<i>Echinochloa crus-galli</i>	2 to 8	12	32
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>	2 to 6	12	32
Brome				
California	<i>Bromus carinatus</i>	2 to 6	12	32
Cheat	<i>Bromus secalinus</i>	2 to 6	12	32
Downy	<i>Bromus tectorum</i>	2 to 6	12	32
Ripgut	<i>Bromus diandrus</i>	2 to 6	12	32
Canarygrass	<i>Phalaris canariensis</i>	1 to 4	12	32
Crabgrass				
Hairy	<i>Digitaria adscendens</i>	2 to 6**	12	32
Large	<i>Digitaria sanguinalis</i>	2 to 6**	12	32
Smooth	<i>Digitaria ischaemum</i>	2 to 6**	12	32
Southern	<i>Digitaria ciliaris</i>	2 to 6**	12	32
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	2 to 6**	12	32
Fall Panicum	<i>Panicum dichotomiflorum</i>	2 to 8	12	32
Field Sandbur	<i>Cenchrus incertus</i>	2 to 6	12	32
Foxtail				
Giant	<i>Setaria faberi</i>	2 to 12	12	32
Green	<i>Setaria viridis</i>	2 to 8	12	32
Yellow	<i>Setaria glauca</i>	2 to 8	12	32
Goosegrass	<i>Eleusine indica</i>	2 to 6**	12	32
Itchgrass	<i>Rottboellia cochlin</i>	2 to 6	12	32
Junglerice	<i>Echinochloa colona</i>	2 to 6	12	32
Lovegrass (Stinkgrass)	<i>Eragrostis cilianensis</i>	2 to 6	12	32
Rabbitsfootgrass	<i>Polypogon monspeliensis</i>	1 to 4	12	32
Red Rice	<i>Oryza sativa</i>	1 to 3	12	32
Rygrass				
Hardy	<i>Lolium remotum</i>	2 to 6	12	32
Italian	<i>Lolium multiflorum</i>	2 to 6	12	32
Seedling Johnsongrass	<i>Sorghum halepense</i>	4 to 10	12	32
Shattercane	<i>Sorghum bicolor</i>	6 to 18	12	32
Southwestern Cupgrass	<i>Eriochloa gracilis</i>	2 to 6	12	32
Sprangletop				
Amazon	<i>Leptochloa panicoides</i>	2 to 6	12	32
Bearded	<i>Leptochloa fascicularis</i>	2 to 6	12	32
Mexican	<i>Leptochloa uninervia</i>	2 to 6	12	32
Red	<i>Leptochloa filiformis</i>	2 to 6	12	32
Texas Panicum	<i>Panicum texanum</i>	2 to 6	12	32
Volunteer Cereals				
Barley	<i>Hordeum vulgare</i>	2 to 6	12	32
Oats	<i>Avena sativa</i>	2 to 6	12	32
Rye	<i>Secale cereale</i>	2 to 6	12	32
Wheat	<i>Triticum aestivum</i>	2 to 6	12	32
Volunteer Corn	<i>Zea mays</i>	4 to 12	12	16
Volunteer Corn	<i>Zea mays</i>	12 to 24	12	32
Volunteer Grain Sorghum	<i>Sorghum bicolor</i>	8 to 12	12	32
Wild Oats	<i>Avena fatua</i>	2 to 6	12	32
Wild Proso Millet	<i>Panicum miliaceum</i>	2 to 10	12	32
Witchgrass	<i>Panicum capillare</i>	2 to 8	12	32
Woolly Cupgrass	<i>Eriochloa villosa</i>	2 to 8	12	32

*Generally occurs between 3-leaf stage and tillering.

**Length of lateral growth.

⁽¹⁾16 fl oz/A (0.121 lb ai/A) = approximately 0.4 fl oz/1000 sq ft

⁽²⁾32 fl oz/A (0.242 lb ai/A) = approximately 0.8 fl oz/1000 sq ft

Add non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt per 50 gals (0.25% v/v).

USE DIRECTIONS FOR ANNUAL BLUEGRASS CONTROL WITH V-10137 1 EC HERBICIDE IN ORNAMENTALS			
GRASS SPECIES	WEED STAGE	APPLICATION RATES	
		MINIMUM RATE fl oz/A	MAXIMUM RATE fl oz/A
Annual Bluegrass (<i>Poa annua</i>)	to 4-leaf	12	32
Apply under favorable soil moisture and humidity that exists within a few days after rainfall or within 7 days after irrigation. Grass needs to be actively growing at time of application(s).			
Apply at weed stage indicated on the label, as reduced control can be expected with more mature annual bluegrass.			
Use the high rate under heavy grass pressure and/or when annual bluegrass is more mature.			
Add non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt per 50 gals (0.25% v/v).			

USE DIRECTIONS FOR PERENNIAL GRASSES IN ORNAMENTALS

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

GRASS SPECIES	WEED HEIGHT (inches)	APPLICATION RATES	
		MINIMUM RATE fl oz/A ⁽¹⁾	MAXIMUM RATE ⁽²⁾ fl oz/A
Bermudagrass (<i>Cynodon dactylon</i>)			
First Application	3 (or up to 6" runners)	12	32
Repeat Application(s) (if regrowth occurs)	3 (or up to 6" runners)	12	32
Foxtail Barley (<i>Hordeum jubatum</i>)			
First Application	2 to 6	12	32
Repeat Application(s) (if regrowth occurs)	2 to 6	12	32
Quackgrass (<i>Elytrigia repens</i>)			
First Application	4 to 8	12	32
Repeat Application(s) (if regrowth occurs)	4 to 8	12	32
Rhizome Johnsongrass (<i>Sorghum halepense</i>)			
First Application	12 to 24	12	32
Repeat Application(s) (if regrowth occurs)	6 to 18	9	16
Wirestem Muhly (<i>Muhlenbergia frondosa</i>)			
First Application	4 to 8	12	32
Repeat Application(s) (if regrowth occurs)	4 to 8	12	32

⁽¹⁾16 fl oz/A (0.121 lb ai/A) = approximately 0.4 fl oz/1000 sq ft

⁽²⁾32 fl oz/A (0.242 lb ai/A) = approximately 0.8 fl oz/1000 sq ft

Add non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt per 50 gals (0.25% v/v).

IMPORTANT

Plant tolerance to V-10137 1 EC Herbicide at labeled rates has been found to be acceptable for the indicated genera and species listed below. Due to variability within species, crop growth stage, environmental conditions, and application techniques, it is recommended that the user determine if the herbicide can be used safely on a few plants prior to widespread application. Neither the seller nor the manufacturer of V-10137 1 EC Herbicide have investigated the safety factor to plants not listed on the label.

NON-BEARING FOOD CROPS

V-10137 1 EC HERBICIDE SHOULD NOT BE APPLIED TO NON-BEARING FRUIT OR NUT CROPS WHICH ARE GROWN FOR ROOT STOCK.

Crop injury to non-bearing fruit and nut crops can occur if V-10137 1 EC Herbicide is improperly applied. V-10137 1 EC Herbicide should not be applied directly over the top of these plant types. Instead spray should be directed at the base of the plant where grassy weeds are growing near the ground.

Non-bearing fruit and nut crops are plants which will not bear fruit or nuts for at least one year following V-10137 1 EC Herbicide application.

COMMON NAME	SCIENTIFIC NAME
Apples	<i>Malus</i> spp.
Berries	<i>Vaccinium</i> spp. <i>Rubus</i> spp.
Cherry, Sweet	<i>Prunus avium</i>
Citrus Fruits	<i>Citrus</i> spp.
Grapes	<i>Vitis</i> spp.
Olives	<i>Olea</i> spp.
Peach	<i>Prunus persica</i>
Pears	<i>Pyrus communis</i>
Prunes	<i>Prunus</i> spp.
Stone Fruits	<i>Prunus</i> spp.
Strawberries	<i>Fragaria</i> spp.
Tree Nuts	
Almond	<i>Prunus triloba</i>
Filbert	<i>Corylus maxima</i>
Pecan	<i>Carya illinoensis</i>
Pistachio	<i>Pistacia vera</i>
Walnut	<i>Juglans</i> spp.

CONIFER TREES

V-10137 1 EC Herbicide can be used to control labeled grasses in Christmas tree farms, conifer nurseries and conifer plantations (but not in forests).

COMMON NAME	SCIENTIFIC NAME
Arborvitae, American	<i>Thuja occidentalis</i>
Cedars	<i>Cedrus</i> spp.
Cypress	<i>Taxodium</i> spp.
Fir, Douglas	<i>Pseudotsuga menziesii</i>
Firs	<i>Abies</i> spp
Hemlock, Canadian/Eastern	<i>Tsuga canadensis</i>
Hemlock, Western	<i>Tsuga heterophylla</i>
Pines	<i>Pinus</i> spp.
Spruces	<i>Picea</i> spp.
Yew	<i>Taxus</i> spp.

NON-CROP OR NON-PLANTED AREAS

The following areas are considered non-crop or non-planted areas: Rights-of-way including railroads, highways, roads, dividers, medians, pipelines, public utility lines, pumping stations, transformer stations and substations. Around airports, electric utilities, commercial buildings, manufacturing plants, storage yards, rail yards, fence lines, parkways, and post-harvest croplands. Also beneath greenhouse benches and around golf courses.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.
Open dumping is prohibited.

PESTICIDE STORAGE

Keep pesticide in original container.
Do not put concentrate or dilute into food or drink containers.
Store in cool, dry place.
Do not store diluted spray.

Emergency Response: For help with any spill, leak, fire or exposure involving this material, call day or night **800-892-0099**.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

[Container language for 2.5 Gallon]

CONTAINER HANDLING

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[Container language for 15 Gallon]

CONTAINER HANDLING

Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[Container language for 120 Gallon]

CONTAINER HANDLING

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

©2016 Valent U.S.A. Corporation

Cobra, *Phoenix* and *Resource* are trademarks and registered trademarks of Valent U.S.A. Corporation
Asana and *Danitol* are registered trademarks of Sumitomo Chemical Company, Ltd.

Harmony, Synchrony and UpBeet are registered trademarks of E.I. du Pont de Nemours and Company

Banvel, Basagran Clearfield, Headline, Pursuit and Raptor are registered trademarks of BASF

Baythroid, Betamix, Betanex, Bronate Advanced, Buctril, Gem, Liberty, Liberty Link and Progress are trademark and registered trademarks of Bayer

Crossbow, Curtail, FirstRate, Frontrow, Grazon, Lorsban, Stinger and Tordon are registered trademarks of Dow AgroSciences LLC

Eminent is a registered trademark of ISAGRO S.p.A.

Flexstar and Warrior are registered trademarks of Syngenta

Orthene is a registered trademark of OMS Investments, Inc. Pounce is a registered trademark of FMC Corporation

Roundup Ready is a registered trademark of Monsanto Company

Storm and Ultra Blazer are registered trademarks of United Phosphorus, Inc.

***Asana* and *Danitol* are restricted use pesticides.**

Manufactured for:

Valent U.S.A. Corporation

P.O. Box 8025

Walnut Creek, CA 94596-8025

www.valent.com

Made in U.S.A.

EPA Reg. No. 59639-132

EPA Est. No.

059639-00132.20160427.V10137.IR4

THE VALENT RETURNABLE KEG

Description: This keg is a closed-system, refillable container designed for easy handling and convenient dispensing of product with no container disposal.

Construction: The keg is made of all stainless steel. Both the gaskets and seals are Viton and are compatible with the Valent product.

Pump System: With the versatility of the keg, either a mechanical pump or an air pressure system may be used to dispense the product.

Coupler: A specific dry-disconnect coupler is required for dispensing product from the keg. This coupler is available through local agricultural equipment suppliers.

Container Capacity: 15 gallons or 56.7 liters (by weight)

ATTENTION!

This is a closed-system container. Do not try to remove the valve from the keg. The coupler required for removal of product is available from local agricultural equipment suppliers. The keg contains tamper evident seals that, if broken, will incur a fee for the user of the keg. Both the coupler and the valve are designed for one-way operation only. Never try to pump any type of material back into the keg.

DIRECTIONS FOR USE

The proper coupler must be attached and engaged before removing any product from the keg. Either a mechanical pump or an air pressure system may be used and connected to the 1-inch NPT thread on the top of the coupler.

IMPORTANT! Attach a hose or pump to the coupler before engaging coupler. This will prevent the user from being splashed in the event that pressure build-up in the keg forces liquid up through the coupler.

To attach and engage the coupler:

1. Pull top of black dust cover back to expose head of valve. The bottom ring of the black dust cover will still be attached to the neck of the valve. Save the dust cover for reuse when returning keg.
2. Before engaging the coupler, securely attach a hose or pump to the threaded connection.
3. Twist coupler onto valve on keg.
and engage coupler by pulling handle straight out to unlock and then pushing handle down into lower position to open internal valve. Handle will automatically lock in place.
4. Secure and engage coupler by pulling handle straight out to unlock and then pushing handle down into lower position to open internal valve. Handle will automatically lock in place.
5. You are now ready to begin the pumping operation.

To remove coupler from container:


1. Release coupler by pulling handle straight out to unlock and then lifting handle into upper position. Handle will automatically lock in place.
2. Lift coupler from keg. As coupler clears top of valve, pull coupler sideways and lift it off the valve.
3. Wipe valve off and replace dust cover.
4. Flush coupler with water.
5. Wipe coupler and store in a clean place.
6. Properly dispose of cleaning towels and rinsate.

RETURNING KEGS

Clean the outside of the keg with water or soap before returning the keg to the distributor. Leave all Valent product labels and stickers securely attached. All Valent product labels, stickers and other information must remain on the keg in order to comply with both State and Federal regulations.

All Valent kegs are tracked using the individual keg serial number stamped in the top of the keg. Distributors are responsible for these kegs that have been assigned to them. Return this keg to the distributor from which it was purchased. Notify the distributor if the keg cannot be returned by the specific time.

Valent U.S.A. Corporation
1600 Riviera Ave., Ste.2 00
Walnut Creek, CA 94596



Supplemental Label



[Bracketed text is optional]

V-10137 1 EC HERBICIDE EPA Reg. No. 59639-132

ACCEPTED

04/29/2016

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 59639-132

APPLICATION OF V-10137 1 EC HERBICIDE IN BERRY LOW GROWING (EXCEPT CRANBERRY AND STRAWBERRY SUBGROUP 13-07G); CANOLA INCLUDING RAPESEED SUBGROUP 20A (EXCEPT FLAX SEED, MUSTARD SEED AND SESAME SEED); POME FRUIT (CROP GROUP 11-10); STEVIA, DRIED LEAVES; STONE FRUIT (CROP GROUP 12-12) AND SUNFLOWER (SUBGROUP 20B)

This supplemental label expires May 2, 2019 and must not be used or distributed after this date.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR V-10137 1 EC HERBICIDE BEFORE APPLYING. USE OF V-10137 1 EC HERBICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR V-10137 1 EC HERBICIDE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Restriction: Do not plant rotational crops until 30 days after application of V-10137 1 EC Herbicide unless the crop is listed on V-10137 1 EC Herbicide label.

CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-10137 1 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
Berry Low Growing (except Cranberry and Strawberry) (Subgroup 13-07G) including: Bearberry Bilberry Blueberry, lowbush Cloudberry Lingonberry Muntries Partridgeberry	45 days	9 to 16 fl oz ⁽⁵⁾	12 to 16 fl oz	Non-ionic Surfactant NIS at 0.25% v/v	None	For repeat applications make a minimum of a 14 day interval. Verify crop tolerance to V-10137 1 EC Herbicide on a small area of the crop, at the desired V-10137 1 EC Herbicide rate and with the same Non-ionic Surfactant (NIS) that will be used on the field. If no crop response is evident seven (7) days after treatment, V-10137 1 EC Herbicide may be used on the entire field at the rate tested and with the same NIS used in the tolerance test. Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.

CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-10137 1 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
<p>Canola Including Rapeseed Subgroup 20A (except Flax Seed, Mustard Seed and Sesame Seed) including (Not for use in California) Borage Crambe Cuphea Echium Gold of Pleasure (Camelina) Hare's Ear Mustard Lesquerella Lunaria Meadowfoam Milkweed Oil Radish Poppy seed Rapeseed (canola) Sweet Rocket</p>	70 days	9 to 12 fl oz ⁽⁵⁾	12 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>Verify crop tolerance to V-10137 1 EC Herbicide on a small area of the crop, at the desired V-10137 1 EC Herbicide rate and with the same Non-ionic Surfactant (NIS) that will be used on the field. If no crop response is evident seven (7) days after treatment, V-10137 1 EC Herbicide may be used on the entire field at the rate tested and with the same NIS used in the tolerance test.</p> <p>Restrictions Do not apply more than 12 fl oz/A (0.091 lb ai/A) per application.</p> <p>Do not apply more than 12 fl oz/A (0.091 lb ai/A) per year.</p> <p>Do not apply after crop has begun bolting. Crop injury may occur when V-10137 1 EC Herbicide is applied during the bloom period.</p>
<p>Pome Fruit (Crop Group 11-10) (including) Apple Azarole Crabapple Loquat Mayhaw Medlar Pear Pear, Asian Quince Quince, Chinese Quince, Japanese Tejocote</p>	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>For repeat applications make on a minimum of a 14 day interval.</p> <p>Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) in a single application.</p> <p>Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.</p>

CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-10137 1 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
Stevia, (dried leaves)	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>For repeat applications make on a minimum of a 14 day interval.</p> <p>V-10137 1 EC Herbicide has not been tested on all varieties. It is the responsibility of the user to test V-10137 1 EC Herbicide on a small portion of the crop to be treated before treating the entire field.</p> <p>Verify crop tolerance to V-10137 1 EC Herbicide on a small area of the crop, at the desired V-10137 1 EC Herbicide rate and with the same Non-ionic Surfactant (NIS) that will be used on the field. If no crop response is evident seven (7) days after treatment, V-10137 1 EC Herbicide may be used on the entire field at the rate tested and with the same NIS used in the tolerance test.</p> <p>Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) in a single application.</p> <p>Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.</p>
<p>Stone Fruit (Crop Group 12-12) including Apricot Apricot, Japanese Capulin Cherry, Black Cherry, Nanking Cherry, Sweet Cherry, Tart; Jujube, Chinese Nectarine Peach Plum Plum, American Plum, Beach Plum, Canada Plum, Cherry Plum, Chickasaw Plum, Damson Plum, Japanese; Plum, Klamath Plum, Prune Plumcot Sloe</p>	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	<p>For repeat applications make on a minimum of a 14 day interval.</p> <p>Restrictions Do not apply more than 16 fl oz/A (0.121 lb ai/A) in a single application.</p> <p>Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.</p>

CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-10137 1 EC HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
Sunflower (Subgroup 20B) including: Calendula Casteroil Plant Chinese Tallowtree Euphorbia Evening Primrose Jojoba Niger Seed Rose Hip Stokes Aster Tallowwood Tea Oil Plant Vernonia	70 days	9 to 16 fl oz ⁽⁵⁾	12 to 32 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v or Crop Oil Concentrate/ Methylated Seed Oil at 1 qt/A or 1% v/v See tank mix label for specific adjuvant recommendations.	2.5 to 4 lb/A	For repeat applications make on a minimum of a 14 day interval. Verify crop tolerance to V-10137 1 EC Herbicide on a small area of the crop, at the desired V-10137 1 EC Herbicide rate and with the same crop oil concentrate that will be used on the field. If no crop response is evident seven (7) days after treatment, V-10137 1 EC Herbicide may be used on the entire field at the rate tested and with the same crop oil used in the tolerance test. Restrictions Do not apply more than 32 fl oz/A (0.242 lb ai/A) per application. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.

⁽¹⁾ V-10137 1 EC Herbicide is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided.

⁽²⁾ See annual and perennial grass control tables for specific use rate recommendations.

⁽³⁾ Non-ionic surfactant (NIS) in this case refers to an adjuvant containing at least 80% non-ionic surfactant. Crop oil concentrate in this case refers to both crop oil concentrate and crop oil concentrate blends. **Acceptable crop oil concentrates** would be those that contain a minimum of 80% oils and 15% emulsifier. **Acceptable crop oil concentrate blends** would be those that contain a minimum of 60% oils and 25-40% surfactants and emulsifiers. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils.

⁽⁴⁾ Use spray grade ammonium sulfate. The use of ammonium sulfate does not take the place of the required adjuvant.

⁽⁵⁾ See DIRECTIONS FOR REDUCED RATE USE TO CONTROL SMALL ANNUAL GRASSES table.

USE DIRECTIONS FOR ANNUAL GRASSES ALL CROPS

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

Restrictions

- Do not exceed the maximum per application rate listed in CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-10137 1 EC HERBICIDE.
- Do not exceed the maximum yearly rate listed in Table 1, CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-10137 1 EC HERBICIDE.

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (inches)	APPLICATION RATES	
			MINIMUM RATE fl oz/A	MAXIMUM RATE ⁽¹⁾ fl oz/A
Barnyardgrass	<i>Echinochloa crus-galli</i>	2 to 8	9	16
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>	2 to 6	9	16
Brome				
California	<i>Bromus carinatus</i>	2 to 6	9	16
Cheat	<i>Bromus secalinus</i>	2 to 6	9	16
Downy	<i>Bromus tectorum</i>	2 to 6	9	16
Ripgut	<i>Bromus diandrus</i>	2 to 6	9	16
Canarygrass	<i>Phalaris canariensis</i>	1 to 4	9	16
Crabgrass				
Hairy	<i>Digitaria adscendens</i>	2 to 6**	9	16
Large	<i>Digitaria sanguinalis</i>	2 to 6**	9	16
Smooth	<i>Digitaria ischaemum</i>	2 to 6**	9	16
Southern	<i>Digitaria ciliaris</i>	2 to 6**	9	16
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	2 to 6**	9	16
Fall Panicum	<i>Panicum dichotomiflor</i>	2 to 8	9	16
Field Sandbur	<i>Cenchrus incertus</i>	2 to 6	9	16
Foxtail				
Giant	<i>Setaria faberi</i>	2 to 12	9	16
Green	<i>Setaria viridis</i>	2 to 8	9	16
Yellow	<i>Setaria glauca</i>	2 to 8	9	16
Goosegrass	<i>Eleusine indica</i>	2 to 6**	9	16
Itchgrass	<i>Rottboellia cochinchinensis</i>	2 to 6	9	16
Junglerice	<i>Echinochloa colona</i>	2 to 6	9	16
Lovegrass (Stinkgrass)	<i>Eragrostis cilianensis</i>	2 to 6	9	16
Rabbitsfootgrass	<i>Polypogon monspeliensis</i>	1 to 4	9	16
Red Rice	<i>Oryza sativa</i>	1 to 3	9	16
Rygrass				
Hardy	<i>Lolium remotum</i>	2 to 6	9	16
Italian	<i>Lolium multiflorum</i>	2 to 6	9	16
Seedling Johnsongrass	<i>Sorghum halepense</i>	4 to 10	9	16
Shattercane	<i>Sorghum bicolor</i>	6 to 18	9	16
Southwestern Cupgrass	<i>Eriochloa gracilis</i>	2 to 6	9	16
Sprangletop				
Amazon	<i>Leptochloa panicoides</i>	2 to 6	9	16
Bearded	<i>Leptochloa fascicularis</i>	2 to 6	9	16
Mexican	<i>Leptochloa uninervia</i>	2 to 6	9	16
Red	<i>Leptochloa filiformis</i>	2 to 6	9	16

continued

USE DIRECTIONS FOR ANNUAL GRASSES
ALL CROPS (continued)

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (inches)	APPLICATION RATES	
			MINIMUM RATE fl oz/A	MAXIMUM RATE ⁽¹⁾ fl oz/A
Texas Panicum	<i>Panicum texanum</i>	2 to 6	9	16
Volunteer Cereals ⁽³⁾				
Barley	<i>Hordeum vulgare</i>	2 to 6	9	16
Oats	<i>Avena sativa</i>	2 to 6	9	16
Rye	<i>Secale cereale</i>	2 to 6	9	16
Wheat ⁽²⁾	<i>Triticum aestivum</i>	2 to 6	9 ⁽²⁾	16
Volunteer Corn ^(2,3)	<i>Zea mays</i>	up to 12	6	12
Volunteer Corn ⁽³⁾	<i>Zea mays</i>	up to 24	9	14
Volunteer Corn ^(2,3)	<i>Zea mays</i>	up to 36	12	16
Volunteer Grain	<i>Sorghum bicolor</i>	8 to 12	9	16
Sorghum				
Wild Oats	<i>Avena fatua</i>	2 to 6	9	16
Wild Proso Millet	<i>Panicum miliaceum</i>	2 to 10	9	16
Witchgrass	<i>Panicum capillare</i>	2 to 8	9	16
Woolly Cupgrass	<i>Eriochloa villosa</i>	2 to 8	9	16

*Generally occurs between 3-leaf stage and tillering.

**Length of lateral growth.

⁽¹⁾ Rates higher than 16 fl oz/A may be applied in certain geographic areas, cropping situations or environmental conditions, where experience has shown that higher rates are needed for satisfactory control of annual grasses. In these situations, rates from 16 to 32 fl oz/A may be applied. **Restriction:** Do not apply more than 12 fl oz/A of V-10137 1 EC Herbicide per application to canola including rapeseed subgroup 20A (except flax seed, mustard seed and sesame seed).

⁽²⁾ When a cereal grain crop (such as wheat) is interseeded for crop establishment or is planted as wind breaks to aid crop establishment, the minimum V-10137 1 EC Herbicide use rate for control is 12 fl oz/A.

⁽³⁾ Includes Roundup Ready®, Liberty Link® and Clearfield® volunteer corn; however not Sethoxydim Resistant volunteer corn.

**DIRECTIONS FOR REDUCED RATE USE TO CONTROL SMALL ANNUAL GRASSES
CANOLA INCLUDING RAPESEED SUBGROUP 20A
(EXCEPT FLAX SEED, MUSTARD SEED AND SESAME SEED)
(REDUCED RATE DIRECTIONS NOT FOR USE IN CALIFORNIA)**

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Regrowth by tillering may occur if application is made when plants are stressed by lack of moisture, excessive moisture, low or high temperatures and/or under very low humidity.

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT (inches)	RATE FL OZ/ ACRE⁽¹⁾
Barnyardgrass	<i>Echinochloa crus-galli</i>	1 to 4	6
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>	1 to 4	8
Crabgrass			
Large	<i>Digitaria sanguinalis</i>	1 to 3*	6
Large	<i>Digitaria sanguinalis</i>	1 to 4*	8
Smooth	<i>Digitaria ischaemum</i>	1 to 3*	6
Smooth	<i>Digitaria ischaemum</i>	1 to 4*	8
Southern	<i>Digitaria ciliar</i>	1 to 4*	8
Fall Panicum	<i>Panicum dichotomiflorum</i>	1 to 4	6
Foxtail			
Giant	<i>Setaria faberi</i>	1 to 4	6
Green	<i>Setaria viridis</i>	1 to 4	6
Millet	<i>Setaria italica</i>	1 to 4	8
Yellow	<i>Setaria glauca</i>	1 to 4	6
Seedling Johnsongrass	<i>Sorghum halepense</i>	1 to 6	8
Shattercane	<i>Sorghum bicolor</i>	4 to 10	6
Texas Panicum	<i>Panicum texanum</i>	1 to 4	8
Volunteer Cereals			
Barley	<i>Hordeum vulgare</i>	1 to 4	8
Oats	<i>Avena sativa</i>	1 to 4	8
Wheat	<i>Triticum aestivum</i>	1 to 4	8
Volunteer Corn**	<i>Zea mays</i>	4 to 12	6
Wild Proso Millet	<i>Panicum miliaceum</i>	1 to 6	6
Wild Oats	<i>Avena fatua</i>	1 to 4	8

*Length of lateral growth

**Not S.R. Corn

⁽¹⁾Always add a non-ionic surfactant at 0.25% v/v total spray volume unless crop specific restrictions and limitations advise otherwise.

**USE DIRECTIONS FOR PERENNIAL GRASSES
(ALL CROPS)**

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

Restrictions

- Do not exceed the maximum per application rate listed in CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-10137 1 EC Herbicide.
- Do not exceed the maximum yearly rate listed in Table 1, CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR V-10137 1 EC HERBICIDE.

GRASS SPECIES	WEED HEIGHT (inches)	APPLICATION RATE	
		MINIMUM RATE fl oz/A	MAXIMUM RATE fl oz/A
Bermudagrass (<i>Cynodon dactylon</i>) First Application	3 (or up to 6" runners)	12	32
Repeat Application(s) (if regrowth occurs)	3 (or up to 6" runners)	12	32
Fescue, Tall (<i>Festuca arundinacea</i>) First Application	4 to 8	12	32
Repeat Application(s) (if regrowth occurs)	4 to 8	12	32
Foxtail Barley (<i>Hordeum jubatum</i>) First Application	2 to 6	12	32
Repeat Application (if regrowth occurs)	2 to 6	12	32
Orchardgrass (<i>Dactylis glomerata</i>) First Application	4 to 8	12	32
Repeat Application(s) (if regrowth occurs)	4 to 8	12	32
Quackgrass (<i>Elytrigia repens</i>) First Application	4 to 12	12	32
Repeat Application(s) (if regrowth occurs)	4 to 12	12	32
Rhizome Johnsongrass (<i>Sorghum halepense</i>) First Application	12 to 24	12	32
Repeat Application(s) (if regrowth occurs)	6 to 18	9	24
Wirestem Muhly (<i>Muhlenbergia frondosa</i>) First Application	4 to 8	12	32
Repeat Application(s) (if regrowth occurs)	4 to 8	12	32
Perennial Bluegrass [Roughstalk (<i>Poa trivialis</i>)] [Kentucky (<i>Poa prantensis</i>)] First Application	2 to 4	12	32
Repeat Application(s)	2 to 4	12	32
Bentgrass (<i>Agrostis</i> spp.) First Application	2 to 4	-	32
Repeat Application(s) (if regrowth occurs)	2 to 4	-	32

TANK MIX - LABEL INFORMATION

The labels for each of the herbicides recommended for tank mixing with V-10137 1 EC Herbicide are unique to the characteristics of those products and contain restrictions and limitations that may be more restrictive than the V-10137 1 EC Herbicide label in certain considerations. Those concerns may include, but are not limited to:

1. Geographic restrictions - all products are not registered for use in all areas and rates may vary from one region of labeled use to another;
2. Crop rotation restrictions;
3. Applicator certification requirements;
4. Worker safety rules (e.g. protective clothing, reentry time, posting);
5. Soil type or soil characteristics (e.g. pH, OM);
6. Maximum dosage or number of applications per season;
7. Rain free period required; or
8. Application timing (e.g. pre-harvest interval)
9. **Restriction:** Do not exceed the total season rates.

THE MOST RESTRICTIVE LABELING OF ANY PRODUCT USED IN A TANK MIX MUST BE FOLLOWED.

TANK MIX APPLICATION OF V-10137 1 EC HERBICIDE AND BROADLEAF HERBICIDES FOR CONTROL OF GRASSES AND BROADLEAF WEEDS

- Apply only to actively growing grass and broadleaf weeds at recommended height or growth stage listed on each label.
- Apply when the first grass or broadleaf weed species in a mixed population reaches the recommended height or growth stage for treatment.
- Apply under favorable soil moisture and humidity that exist a few days after rainfall or within seven days after irrigation.
- Always add the appropriate adjuvant to the spray mix at the rate recommended for each specific tank mix combination.
- Tank mix applications may sometimes result in reduced grass control and possible increases in crop injury as compared to either product used alone. If regrowth occurs, or an additional flush of new grass emerges, make a second application of V-10137 1 EC Herbicide, as specified in the respective size and rate tables.
- **Restriction:** Do not tank mix V-10137 1 EC Herbicide when broadleaf weeds are tall and/or dense enough to prevent proper grass coverage.

MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, add the correct amount of V-10137 1 EC Herbicide. Agitation should create a rippling or rolling action on the water surface.
3. If tank mixing V-10137 1 EC Herbicide with other labeled herbicides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions. Prepare no more spray mixture than is required for the immediate spray operation.
4. Add any required adjuvants (crop oil concentrate, non-ionic surfactant and/or nitrogen solution).
5. Fill spray tank to desired level with water. **Agitation should continue until all spray solution has been applied.**

Failure to agitate the spray solution may result in improper mixing of the herbicides and unsatisfactory weed control. Mixing and compatibility qualities should be verified by a jar test.

INFORMATION ON ANTAGONISM

Tank mixes of V-10137 1 EC Herbicide with postemergence broadleaf herbicides have shown some reduction or failure to control certain grass species which would have otherwise been controlled when V-10137 1 EC Herbicide is applied alone. Activity of the postemergence broadleaf herbicide in the tank mix is not affected.

USE DIRECTIONS FOR POME FRUIT AND STONE FRUIT

Apply V-10137 1 EC Herbicide postemergence to actively growing grasses according to rate table recommendations. Crop injury to pome fruit and stone fruit can occur if V-10137 1 EC Herbicide is improperly applied. V-10137 1 EC Herbicide should not be applied directly over the top of these plant types. Instead spray should be directed at the base of the plant where grassy weeds are growing near the ground.

Restriction: Do not apply V-10137 1 EC Herbicide to pome fruit or stone fruit grown for root stock.

Non-bearing fruit and nut crops are plants which will not bear fruit or nuts for at least one year following V-10137 1 EC Herbicide application.

CANOLA (EXCEPT FLAX)

[Rapeseed Subgroup 20A (except flax seed, mustard seed and sesame seed)]
REDUCED RATE V-10137 1 EC HERBICIDE TANK MIXES WITH BROADLEAF HERBICIDES
(Refer to the use directions tables above for specific grasses and growth stages.)

PRODUCT	APPLICATION RATES/ACRE	ADJUVANT RECOMMENDATIONS	AMMONIUM SULFATE	
	ANNUAL GRASSES ⁽¹⁾		GROUND	AIR
V-10137 1 EC ⁽²⁾ Herbicide + Liberty ^{®(3)}	8 to 10 fl oz + 28 to 34 fl oz	NIS at 0.25% v/v	3 lb/A	3 lb/A
V-10137 1 EC ⁽²⁾ Herbicide + Stinger ^{®(4)}	8 to 10 fl oz + 0.33 pts/A	NIS at 0.25% v/v	3 lb/A	3 lb/A

⁽¹⁾ Annual grasses and sizes controlled with these tank mixtures are those that are identified in the DIRECTIONS FOR REDUCED RATE USE TO CONTROL SMALL ANNUAL GRASSES table.

⁽²⁾ **Restriction:** Do not apply V-10137 1 EC Herbicide tank mix during or after bolting or flowering or crop injury will occur.

⁽³⁾ For use only on Liberty Link Canola.

⁽⁴⁾ See Stinger label for weeds controlled.

TANK MIX APPLICATION OF V-10137 1 EC HERBICIDE AND INSECTICIDES FOR CONTROL OF GRASS WEEDS AND INSECTS IN SUNFLOWER

PRODUCT ⁽¹⁾	APPLICATION RATES/ACRE ⁽²⁾		ADJUVANT RECOMMENDATION
	ANNUAL GRASSES	PERENNIAL GRASSES	
V-10137 1 EC Herbicide + Asana [®] XL	9 to 12 fl oz + Refer to Asana XL label	12 to 24 fl oz + Refer to Asana XL label	NIS at 0.25% v/v + AMS at 2.5 lb/A
V-10137 1 EC Herbicide + Warrior [®]	9 to 12 fl oz + Refer to Warrior label	12 to 24 fl oz + Refer to Warrior label	NIS at 0.25% v/v + AMS at 2.5 lb/A

⁽¹⁾ Refer to V-10137 1 EC Herbicide and insecticide label for rates and weeds and insects controlled.

⁽²⁾ If grass regrowth occurs, or an additional flush of new grass emerges, make a second application of V-10137 1 EC Herbicide alone (without a tank mix insecticide) according to the appropriate size and rate recommendations.

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

©2016 Valent U.S.A. Corporation

Asana is a registered trademark of Sumitomo Chemical Company, Ltd.
Clearfield is a registered trademark of BASF
Liberty and Liberty Link are registered trademarks of Bayer
Roundup Ready is a registered trademark of Monsanto Company
Stinger is a registered trademark of Dow AgroSciences LLC
Warrior is a registered trademark of Syngenta

Manufactured for:

Valent U.S.A. Corporation

P.O. Box 8025

Walnut Creek, CA 94596-8025

Made in U.S.A.

059639-00003.20160407.V10137_multicrops_supp